arXiv:1401.6852v1 [math.OA] 27 Jan 2014

CROSSED PRODUCT BY ACTIONS OF FINITE GROUPS WITH THE ROKHLIN PROPERTY

LUIS SANTIAGO

Abstract. We introduce and study a Rokhlin-type property for actions of finite groups on (not necessarily unital) C*-algebras. We show that the corresponding crossed product C*-algebras can be locally approximated by C*-algebras that are stably isomorphic to closed two-sided ideals of the given algebra. We then use this result to prove that several classes of C*-algebras are closed under crossed products by finite group actions with this Rokhlin property.

Contents

1. Introduction 1
2. Preliminary definitions and results 2
3. The Rokhlin property 5
4. Local approximation of $A \rtimes_\alpha G$ 13
5. Permanence properties 16
References 22

1. Introduction

The Rokhlin property is a form of freeness for group actions on noncommutative C*-algebras. For finite group actions on unital C*-algebras this property first appeared (under different names) in [20] and [12] for actions of $\mathbb{Z}_p$ on UHF-algebras, and in [15] for actions of finite groups on arbitrary unital C*-algebras. Let us recall the definition of the Rokhlin property:

Definition 1. ([17, Definition 3.1]) Let A be a unital C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A. The action α is said to have the *Rokhlin property* if there exists a partition of unity $(p_g)_{g\in G} \subset A_\infty$ (see Subsection 2.1 for the definition of A_∞) consisting of projections such that

$$\alpha_g(p_h) = p_{gh},$$

for all $g, h \in G$.

When A is separable this property can be reformulated in terms of finite subsets and projections of A ([28, Definition 1.1]). The Rokhlin property is quite restrictive and is not useful if there is no abundance of projections. For instance, if either the K_0-group or the K_1-group of the algebra is isomorphic to $\mathbb{Z}$ then the algebra does not have any action of a finite group with the Rokhlin porperty (this follows by [17, Theorem 3.13]). On the contrast, if a C*-algebra absorbs tensorially the UHF-algebra of infinite type M_{n^∞} and G is a finite group of order n then the algebra has an action of G with the Rokhlin property. In fact, in this case the actions of G with the Rokhlin property are generic ([29, Theorem 5.17]).

Date: January 28, 2014.

For C*-algebras that are not unital there are at least two Rokhlin-type properties. One of them is called the multiplier Rokhlin property ([27, Definition 2.15]), and is defined using projections of the multiplier algebra of the given C*-algebra. The other one, which is simply called the Rokhlin property ([22, Definition 3.1]), is defined for σ-unital C*-algebras using projections in the central sequence algebra of the given C*-algebra. In this paper we introduce a Rokhlin-type property (Definition 2) which somehow resembles [28, Definition 1.1]. It is defined using positive elements of the algebra instead of projections. This new Rokhlin-type property is essentially that of [22, Definition 3.1] (they agree for separable C*-algebras), but it has the advantage that it can be defined for general C*-algebras (not just for σ-unital ones). Actions with the multiplier Rokhlin property or the Rokhlin property in the sense of [22, Definition 3.1] also have this new Rokhlin property. Also, it can be seen that if the algebra is unital our definition agrees with the standard Rokhlin property for unital C*-algebras. In this paper we also study the crossed product C*-algebras obtained by actions of finite groups with the Rokhlin property on different classes of C*-algebras. Our main motivation comes from the study of crossed products by finite group actions on stably projectionless C*-algebras and our objective is to extend the results of [23] to non-unital C*-algebras. The problem of classifying actions with the Rokhlin property will be considered in [13].

The paper is organized as follows. In Section 3 we define and study a Rokhlin-type property for actions of finite groups on arbitrary C*-algebras. Section 4 is dedicated to show that the crossed product C*-algebra by an action of a finite group on a C*-algebra with this Rokhlin property can be locally approximated by matrix algebras over hereditary C*-subalgebras of the given C*-algebra (Theorem 2). This result may be considered as a nonunital version of [23, Theorem 3.2]. In Section 5 we use this approximation result to show that several classes of C*-algebras are closed under crossed products by finite group actions with the Rokhlin property (Theorem 3). These classes of C*-algebras include purely infinite C*-algebras, C*-algebras of stable rank one, C*-algebras of real rank zero, C*-algebras of finite nuclear dimension, separable $\mathcal{D}$-stable C*-algebras, where $\mathcal{D}$ is a strongly self-absorbing C*-algebra, simple C*-algebras, simple C*-algebras with strict comparison of positive elements, simple stably projectionless C*-algebras, and C*-algebras that are stably isomorphic to sequential direct limits of one-dimensional NCCW-complexes.

I would like to thank Eusebio Gardella for useful discussions concerning the subject matter of this paper.

2. Preliminary definitions and results

2.1. **Central sequence algebras.** Let A be a C*-algebra. Let us denote by A^∞ the quotient of $\ell_\infty(\mathbb{N}, A)$ by the ideal $c_0(\mathbb{N}, A)$. That is,

$$A^\infty := \ell_\infty(\mathbb{N}, A)/c_0(\mathbb{N}, A).$$

The elements of $\ell_\infty(\mathbb{N}, A)$ will be written as sequences of elements of A. If $(a_n)_n \in \ell_\infty(\mathbb{N}, A)$ then its image in the quotient A^∞ will be denoted by $[(a_n)_n]$.

Let B be a C*-subalgebra of A. Then we will identify B with its image in A^∞ by the embedding

$$a \in B \longmapsto (a, a, \cdots) \in A^\infty.$$

Similarly, we will identify $B^\infty \cap B'$ with the corresponding C*-subalgebra of $A^\infty \cap B'$. Let us denote by A_∞ and $\mathrm{Ann}(B, A^\infty)$ the C*-algebras

$$A_\infty := A^\infty \cup A', \quad \mathrm{Ann}(B, A^\infty) := \{a \in A^\infty \cap B' : ab = 0, \ \ \forall b \in B\}.$$

Then $\mathrm{Ann}(B, A^\infty)$ is a closed two-sided ideal of $A^\infty \cap B'$. Denote the corresponding quotient by $\mathrm{F}(B, A)$ and the quotient map by π. We call $\mathrm{F}(A) := \mathrm{F}(A, A)$ the *central sequence algebra* of A. Note that if B is σ-unital then $\mathrm{F}(B, A)$ is unital. Let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on the C*-algebra A and let B be a G-invariant subalgebra of A. Then α induces natural actions on A^∞, A_∞, $\mathrm{F}(A)$, and $\mathrm{F}(B, A)$. For simplicity we will denote all these actions again by α.

Lemma 1. Let A be a C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A. Let J be a G-invariant ideal. Then J has a quasi-central approximate unit consisting of G-invariant positive contractions.

Proof. By [24, Theorem 3.12.14] it is enough to show that J has a G-invariant approximate unit. Let $(u_i)_{i\in I}$ be an approximate unit for J. For each $i \in I$ put $v_i = \frac{1}{|G|}\sum_{g\in G} \alpha_g(u_i)$. Then it is clear that $\|v_i\| \leq 1$ for all $i \in I$, $v_i \leq v_j$ for all $i, j \in I$ with $i \leq j$, and $\alpha_g(v_i) = v_i$ for all $i \in I$ and $g \in G$. Let $a \in J$ and let $\epsilon > 0$. Then there exists i_0 such that

$$\|u_i\alpha_{g^{-1}}(a) - \alpha_{g^{-1}}(a)\| < \epsilon,$$

for all $i \geq i_0$ and $g \in G$. Now using that *-automorphisms are norm preserving we get

$$\|\alpha_g(u_i)a - a\| = \|\alpha_g(u_i\alpha_{g^{-1}}(a) - \alpha_{g^{-1}}(a))\| = \|u_i\alpha_{g^{-1}}(a) - \alpha_{g^{-1}}(a)\| < \epsilon,$$

for all $i \geq i_0$ and $g \in G$. Hence, by the triangle inequality

$$\|v_ia - a\| \leq \frac{1}{|G|}\sum_{g\in G} \|\alpha_g(u_i)a - a\| < \frac{1}{|G|}(|G|\epsilon) = \epsilon,$$

for all $i \geq i_0$. This shows that $(v_i)_{i\in I}$ is an approximate unit for J. □

Lemma 2. Let A be a C*-algebra, let $\alpha\colon G \to A$ be an action of a finite group G on A, and let B be a σ-unital C*-subalgebra of A. Let F_1 be a finite subset of $A^\infty \cap B'$, let F_2 be a finite subset of $\mathrm{Ann}(B, A^\infty)$, and let F_3 be a finite subset of

$$\{a \in A^\infty : ab = b \text{ for all } b \in B\}.$$

Then there exists a positive contraction d in the fixed point algebra $(B^\infty \cap B')^G$ (considered as a subalgebra of $(A^\infty \cap B')^G$) such that

$$\begin{aligned} db &= b, && \forall b \in B,\\ da &= ad, && \forall a \in F_1,\\ da &= 0, && \forall a \in F_2,\\ da &= a, && \forall a \in F_3. \end{aligned}$$

Proof. Let $(u_n)_{n\in\mathbb{N}} \subset B^G$ be a countable approximate unit for B (this approximate unit exists by the previous lemma). Let F_1, F_2, and F_3 be as in the statement of the lemma. Without loss of generality we may assume that their cardinalities is the same and that their elements are positive. Write

$$\begin{aligned} F_1 &= \{a^{(i)} = [(a_n^{(i)})_n] : 1 \leq i \leq m\},\\ F_2 &= \{b^{(i)} = [(b_n^{(i)})_n] : 1 \leq i \leq m\},\\ F_3 &= \{c^{(i)} = [(c_n^{(i)})_n] : 1 \leq i \leq m\}. \end{aligned}$$

Then for each $j \in \mathbb{N}$ there exists $k_j \in \mathbb{N}$ such that

$$\|u_ja_n^{(i)} - a_n^{(i)}u_j\| < \frac{1}{j}, \qquad \|u_jb_n^{(i)}\| < \frac{1}{j}, \qquad \|u_jc_n^{(i)} - c_n^{(i)}\| < \frac{1}{j}, \tag{1}$$

for $n \geq k_j$ and $1 \leq i \leq m$. Moreover, we may choose the sequence $(k_j)_{j\in\mathbb{N}}$ satisfying $k_j < k_{j+1}$ for all $j \in \mathbb{N}$. Define $d = [(d_n)_n] \in B^\infty \subseteq A^\infty$ by

$$d_n = \begin{cases} u_1 & \text{for } 1 \leq n < k_2\\ u_j & \text{for } k_{j+1} \leq n < k_{j+2}. \end{cases}$$

Then $d \in B^\infty \cap B'$ and $db = b$ for all $b \in B$, since $(u_n)_{n\in\mathbb{N}}$ is an approximate unit for B. In addition, by the inequalities in (1) we have

$$da^{(i)} = a^{(i)}, \qquad db^{(i)} = 0, \qquad dc^{(i)} = c^{(i)},$$

for $1 \leq i \leq m$. This concludes the proof of the lemma. □

2.2. **The Cuntz semigroup.** Let us briefly recall the definition of the Cuntz semigroup of a C*-algebra. Let A be a C*-algebra and let $a, b \in A$ be positive elements. We say that a is *Cuntz subequivalent* to b, and denote this by $a \precsim b$, if there exists a sequence $(d_n)_{n\in\mathbb{N}} \subset A$ such that $d_n^* b d_n \to a$ as $n \to \infty$. We say that a is *Cuntz equivalent* to b, and denote this by $a \sim b$, if $a \precsim b$ and $b \precsim a$. It is easy to see that $\precsim$ is a preorder relation on A_+ and that $\sim$ is an equivalence relation. We denote the Cuntz equivalence class of an element $a \in A_+$ by $[a]$. The *Cuntz semigroup* of A, denoted by $\mathrm{Cu}(A)$, is the set of Cuntz equivalence classes of positive elements of $A \otimes \mathcal{K}$, where $\mathcal{K}$ denotes the algebra of compact operators on a separable Hilbert space. The addition on $\mathrm{Cu}(A)$ is given by $[a] + [b] := [a' + b']$, where $a', b' \in (A \otimes \mathcal{K})_+$ are such that $a \sim a'$, $b \sim b'$, and $a'b' = 0$. The preorder relation $\precsim$ induces an order relation on $\mathrm{Cu}(A)$; that is, $[a] \leq [b]$ if $a \precsim b$. We say that $[a]$ is *compactly contained* in $[b]$, and denote this by $[a] \ll [b]$, if whenever $[b] \leq \sup_n [b_n]$, for some increasing sequence $([b_n])_{n\in\mathbb{N}}$, one has $[a] \leq [b_k]$ for some k. A sequence $([a_n])_{n\in\mathbb{N}}$ is said to be rapidly increasing if $[a_n] \ll [a_{n+1}]$ for all n. It was shown in [8] that $\mathrm{Cu}(A)$ is closed under taking suprema of increasing sequences and that every element of $\mathrm{Cu}(A)$ is the supremum of a rapidly increasing sequence. In particular, we have

$$[(a-\epsilon)_+] \ll [a], \quad [a] = \sup_{\epsilon>0}[(a-\epsilon)_+],$$

for all $a \in (A \otimes \mathcal{K})_+$ and all $\epsilon > 0$. Here $(a-\epsilon)_+$ denotes the element obtained by evaluating—using functional calculus—the continuous function $f(t) = \max(0, t-\epsilon)$, with $t \in (0, \infty)$, at the element a.

The following lemma states two well known result regarding Cuntz subequivalence. The first statement was proved in [19, Lemma 2.2] in the case $\delta = 0$ and in [32, Lemma 1] for an arbitrary δ. The second statement was shown in [19, Lemma 2.3].

Lemma 3. Let A be a C*-algebra and let $a, b \in A_+$. The following statements hold:

(i) If $\|a - b\| < \epsilon$, then $(a - (\epsilon + \delta))_+ \precsim (b - \delta)_+$ for all $\delta > 0$;
(ii) If $a \precsim b$ then for every $\epsilon > 0$ there exists $\delta > 0$ such that $(a - \epsilon)_+ \precsim (b - \delta)_+$.

We say that $\mathrm{Cu}(A)$ is *almost unperforated* if for any $[a], [b] \in \mathrm{Cu}(A)$ satisfying $(n+1)[a] \leq n[b]$, for some $n \in \mathbb{N}$, one has $[a] \leq [b]$. Consider the set $\mathrm{L}(A)$ of all functionals $\lambda \colon \mathrm{Cu}(A) \to [0, \infty]$ that are additive, order-preserving, and that preserve suprema of increasing sequences. We say that a simple C*-algebra A has *strict comparison of positive elements* if for $[a], [b] \in \mathrm{Cu}(A)$, $\lambda([a]) < \lambda([b])$ for all $\lambda \in \mathrm{L}(A)$ satisfying $\lambda([b]) < \infty$, implies $[a] \leq [b]$. The following is [39, Lemma 6.1].

Lemma 4. Let A be a simple C*-algebra. Then $\mathrm{Cu}(A)$ is almost unperforated if and only if A has strict comparison of positive elements.

Lemma 5. Let A be a C*-algebra and let I be a σ-unital closed two-sided ideal of A. Suppose that $\mathrm{Cu}(A)$ is almost unperforated, then $\mathrm{Cu}(A/I)$ is almost unperforated.

Proof. Let $\pi \colon A \to A/I$ denote the quotient map. Let $n \in \mathbb{N}$ and $x, y \in \mathrm{Cu}(A/I)$ be such that

$$(n+1)x \leq ny.$$

Choose $a, b \in (A \otimes \mathcal{K})_+$ such that $[\pi(a)] = x$ and $[\pi(b)] = y$. Then it follows by [6, Theorem 1.1] that,

$$(n+1)[a] + z \leq n[b] + z,$$

where z is the largest element of $\mathrm{Cu}(I)$ (this element exists because I is σ-unital). Since $2z = z$ we have

$$(n+1)([a]+z) \leq n([b]+[z]).$$

Using now that $\mathrm{Cu}(A)$ is almost unperforated we get $[a]+z \leq [b]+z$. Hence,

$$x = \mathrm{Cu}(\pi)([a]+z) \leq \mathrm{Cu}(\pi)([b]+z) = y.$$

This implies that $\mathrm{Cu}(A/I)$ is almost unperforated. □

2.3. **Strongly self-absorbing C*-algebras.** Recall that a unital nuclear C*-algebra $\mathcal{D}$ is said to be *strongly self-absorbing* if there exists a *-isomorphism $\phi\colon \mathcal{D} \to \mathcal{D} \otimes \mathcal{D}$ that is approximately unitarily equivalent to the map $i\colon \mathcal{D} \to \mathcal{D} \otimes \mathcal{D}$ given by $i(a) = a \otimes 1_{\mathcal{D}}$ for all $a \in \mathcal{D}$. A C*-algebra A is $\mathcal{D}$-stable if $A \otimes \mathcal{D} \cong A$.

The following result was proved in [16, Proposition 4.1] under the assumption that the algebra $\mathcal{D}$ is K_1-injective. This assumption is in fact redundant since every strongly self-absorbing C*-algebra is K_1-injective ([42, Theorem 3.1 and Remark 3.3]).

Lemma 6. Let A and $\mathcal{D}$ be separable C*-algebras such that $\mathcal{D}$ is strongly self-absorbing. Then A is $\mathcal{D}$-stable if and only if for any $\epsilon > 0$ and any finite subsets $F \subset A$ and $G \subset \mathcal{D}$ there exists a completely positive contractive map $\phi\colon \mathcal{D} \to A$ such that

(i) $\|b\phi(1_{\mathcal{D}}) - b\| < \epsilon$,
(ii) $\|b\phi(d) - \phi(d)b\| < \epsilon$,
(iii) $\|b(\phi(dd') - \phi(d)\phi(d'))\| < \epsilon$,

for all $b \in F$ and $d, d' \in G$.

3. The Rokhlin property

In this section we introduce a Rokhlin-type property for actions of finite groups on not necessarily unital C*-algebras and study several permanence properties of it. We also study its relation with the Multiplier Rokhlin property ([27, Definition 2.15]) and with the Rokhlin property as defined in [22, Definition 3.1].

Definition 2. Let A be a C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A. We say that α has the *Rokhlin property* if for any $\epsilon > 0$ and any finite subset $F \subset A$ there exist mutually orthogonal positive contractions $(r_g)_{g\in G} \subset A$ such that

(i) $\|\alpha_g(r_h) - r_{gh}\| < \epsilon$, for all $g, h \in G$;
(ii) $\|r_g a - a r_g\| < \epsilon$, for all $a \in F$ and $g \in G$;
(iii) $\|(\sum_{g\in G} r_g)a - a\| < \epsilon$, for all $a \in F$.

The elements $(r_g)_{g\in G}$ will be called *Rokhlin elements* for α.

Proposition 1. Let A be a C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A. Then the following are equivalent:

(i) α has the Rokhlin property.
(ii) For any finite subset $F \subset A$ there exist mutually orthogonal positive contractions $(r_g)_{g\in G} \subset A^\infty \cap F'$ such that
  (a) $\alpha_g(r_h) = r_{gh}$, for all $g, h \in G$;
  (b) $(\sum_{g\in G} r_g)b = b$, for all $b \in F$.
(iii) For any separable C*-subalgebra $B \subseteq A$ there are orthogonal positive contractions $(r_g)_{g\in G} \subset A^\infty \cap B'$ such that
  (a) $\alpha_g(r_h) = r_{gh}$, for all $g, h \in G$;
  (b) $(\sum_{g\in G} r_g)b = b$, for all $b \in B$.

(iv) For any separable G-invariant C*-subalgebra $B \subseteq A$ there are projections $(p_g)_{g\in G} \subset \mathrm{F}(B,A)$ such that
   (a) $\alpha_g(p_h) = p_{gh}$, for all $g, h \in G$;
   (b) $\sum_{g\in G} p_g = 1_{\mathrm{F}(B,A)}$.

Proof. (i)⇒(iii). Let B be a separable C*-subalgebra of A. Choose finite subsets $(F_n)_{n\in\mathbb{N}}$ of A such that $F_n \subset F_{n+1}$ for all n, and $\overline{\bigcup_{n\in\mathbb{N}} F_n} = B$. For each n choose Rokhlin elements $(r_g^{(n)})_{g\in G}$ such that

$$\|\alpha_g(r_h^{(n)}) - r_{gh}^{(n)}\| < \frac{1}{n}, \quad \|r_g^{(n)}b - br_g^{(n)}\| < \frac{1}{n}, \quad \|(\sum_{g\in G} r_g^{(n)})b - b\| < \frac{1}{n},$$

for all $g, h \in G$ and all $b \in F_n$. Let $r_g = \left(r_g^{(n)}\right)_n \in A^\infty$. Then $\alpha_g(r_h) = r_{gh}$ for all $g, h \in G$, and

$$r_g b = br_g, \quad (\sum_{g\in G} r_g)b = b,$$

for all $b \in \bigcup_{n\in\mathbb{N}} F_n$. Using that $\bigcup_{n\in\mathbb{N}} F_n$ is dense in B we get the previous equalities for all $b \in B$. This shows that the elements r_g, with $g \in G$, belong to $A^\infty \cap B'$ and that the satisfy (a) and (b) of (iii).

(iii)⇒(ii). This is clear by taking B to be the C*-subalgebra generated by the finite set F.

(ii)⇒(i). Let F be a finite subset of A. Let $(r_g)_{g\in G} \subset A^\infty \cap F'$ be elements satisfying conditions (a) and (b) of (ii). By [21, Lemma 10.1.12] these elements can be lifted to mutually orthogonal positive contractions $((r_{g,n})_n)_{g\in G} \in \ell_\infty(\mathbb{N}, A)$. Let $\epsilon > 0$. Then using the definition of A^∞ and that the elements $(r_g)_{g\in G}$ satisfy conditions (a) and (b) of (ii) we get

$$\begin{aligned}
&\|\alpha_g(r_{h,n}) - r_{gh,n}\| < \epsilon, \qquad \forall g, h \in G,\\
&\|r_{g,n}b - br_{g,n}\| < \epsilon, \qquad \forall g \in G, \forall b \in F,\\
&\|(\sum_{g\in G} r_{g,n})b - b\| < \epsilon, \qquad \forall b \in F,
\end{aligned}$$

for some $n \in \mathbb{N}$. This shows that α has the Rokhlin property.

(iii)⇒(iv). Let B be a separable G-invariant C*-subalgebra of A. By (iii) there exist positive orthogonal contractions $(r_g) \subset A^\infty \cap B'$ satisfying (a) and (b) of (iii). Let $\pi\colon A^\infty \cap B' \to \mathrm{F}(A,B)$ be the quotient map. Set $p_g = \pi(r_g)$. Then by (b) of (iii) we have $\sum_{g\in G} p_g = 1_{\mathrm{F}_{B,A}}$. Also, by (a) of (iii) we get $\alpha_g(p_h) = p_{g,h}$ for all $g, h \in G$. Since the elements $(r_g)_{g\in G}$ are orthogonal, the elements $(p_g)_{g\in G}$ are orthogonal. It follows that $p_h^2 = p_h \sum_{g\in G} p_g = p_h$ for all $h \in G$. Hence, $(p_g)_{g\in G}$ are projections.

(iv)⇒(ii). Let F be a finite subset of A and let B be the C*-subalgebra of A generated by the finite set $\bigcup_{g\in G} \alpha_g(F)$. Then B is G-invariant and separable. Let $(p_g)_{g\in G} \subset \mathrm{F}(B,A)$ be projections satisfying conditions (a) and (b) of (iv). Then by [21, Lemma 10.1.12] there exit mutually orthogonal positive contractions $(f_g)_{g\in G} \subset A^\infty \cap B'$ such that $\pi(f_g) = p_g$. Note that these elements satisfy

$$\alpha_g(f_h)b = f_{gh}b, \quad (\sum_{g\in G} f_g)b = b,$$

for all $g, h \in G$ and all $a \in B$. By Lemma 2 applied to the finite sets $F_1 = \{f_g : g \in G\}$, $F_2 = \{\alpha_g(f_h) - f_{gh} : g, h \in G\}$, and $F_3 = \{\sum_{g\in G} f_g\}$, there exists a positive contraction $d \in (A^\infty \cap B')^G$ such that

$$db = b, \qquad df_g = f_g d, \qquad \alpha_g(f_h)d = f_{gh}d, \qquad (\sum_{g\in G} f_g)d = d,$$

for all $b \in B$, and $g, h \in G$. For each $g \in G$ set $r_g = df_g d$. Then $(r_g)_{g\in G} \subset A^\infty \cap B'$. Also, by the previous equations and that $d \in (A^\infty \cap B')^G$ we get

$$\begin{aligned} &\sum_{g\in G} r_g = d^2, \\ &\alpha_g(r_h) = r_{gh} \qquad \forall g, h \in G, \\ &r_g r_h = 0 \qquad \forall g, h \in G, g \neq h. \end{aligned} \tag{2}$$

Therefore, since $(r_g)_{g\in G} \subset A^\infty \cap F'$ and $d^2 b = b$ for all $b \in B$, it follows that the elements $(r_g)_{g\in G}$ satisfy conditions (a) and (b) of (ii). □

Corollary 1. Let A be a unital C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A. Then α has the standard Rokhlin as in [28, Definition 1.1] (without the assumption of separability) if and only if it has the Rokhlin property as defined in Definition 2.

Proof. The forward implication is clear. Let us show the opposite implication. Let A and α be as in the statement of the corollary. Let F be a finite subset of A and let B be the unital C*-subalgebra of A generated by $\bigcup_{g\in G} \alpha_g(F)$. It is clear that B is separable and G-invariant. Also, $\mathrm{F}(B, A) = A^\infty \cap B'$ since B is unital. By (iv) of Proposition 1 there exists a partition of unity $(p_g)_{g\in G} \subset A^\infty \cap B'$ consisting of projections such that

$$\alpha_g(p_h) = p_{gh},$$

for all $g, h \in G$. Let $\phi\colon \mathbb{C}[G] \to A^\infty \cap B'$ be the *-homomorphism defined by $\phi(\sum_{g\in G} z_g g) = \sum_{g\in G} z_g p_g$, where $z_g \in \mathbb{C}$ for all $g \in G$. Using that the C*-algebra $\mathbb{C}[G] \cong \mathbb{C}^{|G|}$ is semiprojective by [21, Lemma 14.1.5 and Theorem 14.2.1], we can lift ϕ to a unital map from $\mathbb{C}[G]$ to $\ell_\infty(\mathbb{N}, A)$. In other words, there exists a partition of unity $(q_g)_{g\in G} \subset \ell_\infty(\mathbb{N}, A)$, with $q_g = (q_g^{(n)})_{n\in\mathbb{N}}$, consisting of projections such that

$$\begin{aligned} &\sum_{g\in G} q_g^{(n)} = 1, \quad \forall n \in \mathbb{N}, \\ &\lim_{n\to\infty} (\alpha_g(q_h^{(n)}) - q_{g,h}^{(n)}) = 0, \quad \forall g, h \in G, \\ &\lim_{n\to\infty} (q_g^{(n)} a - a q_g^{(n)}) = 0, \quad \forall a \in F. \end{aligned}$$

This implies that α has the Rokhlin property as in [28, Definition 1.1]. □

Let A be a σ-unital C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A. We say that α has the *Rokhlin property in the sense of* [22, Definition 3.1] if there exist projections $(p_g)_{g\in G} \subset \mathrm{F}(A)$ such that

$$\sum_{g\in G} p_g = 1_{\mathrm{F}(A)}, \qquad \alpha_g(p_h) = p_{g,h} \qquad \forall g, h \in G.$$

Corollary 2. Let A be a σ-unital C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A. Suppose that α has the Rokhlin property in the sense of [22, Definition 3.1], then α has the Rokhlin property. Moreover, if A is separable then α has the Rokhlin property in the sense of [22, Definition 3.1] if and only if it has the Rokhlin property.

Proof. This is a consequence of the equivalence of (i) and (iii) of Proposition 1. □

Let A be a C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A. Let $\mathrm{M}(\alpha)$ denote the extension of α to the multiplier algebra $\mathrm{M}(A)$ of A. Recall that α has the *multiplier Rokhlin property* ([27, Definition 2.15]) if for every finite set $F \subset A$ and every $\epsilon > 0$, there are mutually orthogonal projections $(p_g)_{g\in G} \subset \mathrm{M}(A)$ such that:

(i) $\|\mathrm{M}(\alpha)_g(p_h) - p_{gh}\| < \epsilon$ for all $g, h \in G$;
(ii) $\|p_g a - a p_g\| < \epsilon$ for all $g \in G$ and all $a \in F$;
(iii) $\sum_{g\in G} p_g = 1_{\mathrm{M}(A)}$.

Corollary 3. Let A be a C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A. If α has the multiplier Rokhlin property then α has the Rokhlin property.

Proof. Let A and α be as in the statement of the corollary. Let us show that (A, α) satisfies (iv) of Proposition 1. Let B be a separable G-invariant C*-subalgebra of A. Choose finite subsets $(F_n)_{n\in\mathbb{N}}$ of B consisting of contractions such that $\bigcup_{n\in\mathbb{N}} F_n$ is dense in the unit ball of B. For each $n \in \mathbb{N}$ let $(p_g^{(n)})_{g\in G} \subset \mathrm{M}(A)$ be mutually orthogonal projections satisfying properties (i), (ii), and (iii) given above for F_n and $\epsilon = \frac{1}{n}$. Select a G-invariant approximate unit $(u_\lambda)_{\lambda\in\Lambda}$ for A that is quasi-central for $\mathrm{M}(A)$ (this approximate unit exists by Lemma 1). Let $n \in \mathbb{N}$ be fixed. Then for each $a \in F_n$ we have

$$\lim_\lambda \|(u_\lambda p_g^{(n)} u_\lambda)(u_\lambda p_h^{(n)} u_\lambda)a\| = \lim_\lambda \|p_g^{(n)} p_h^{(n)} a u_\lambda^4\| = 0 < \frac{1}{n}, \quad \forall g, h \in G, \ \ g \neq h$$

$$\lim_\lambda \|\alpha_g(u_\lambda p_h^{(n)} u_\lambda)a - u_\lambda p_{gh}^{(n)} u_\lambda a\| = \lim_\lambda \|(\mathrm{M}(\alpha)_g(p_h^{(n)}) - p_{gh}^{(n)}) a u_\lambda^2\| < \frac{1}{n}, \quad \forall g, h \in G,$$

$$\lim_\lambda \|(u_\lambda p_g^{(n)} u_\lambda)a - a(u_\lambda p_g^{(n)} u_\lambda)\| = \lim_\lambda \|(p_g^{(n)} a - a p_g^{(n)}) u_\lambda^2\| < \frac{1}{n}, \quad \forall g \in G,$$

$$\lim_\lambda \|(\sum_{g\in G} u_\lambda p_g^{(n)} u_\lambda)a - a\| = \|(\sum_{g\in G} p_g^{(n)})a - a\| = 0 < \frac{1}{n},$$

Therefore, there exists a positive contraction $u \in A$ such that for $r_g^{(n)} := u p_g^{(n)} u \in A$ the following hold:

$$\|r_g^{(n)} r_h^{(n)} a\| < \frac{1}{n}, \quad \forall g, h \in G, \ \ g \neq h, \quad \forall a \in F_n$$

$$\|(\alpha_g(r_h^{(n)}) - r_{gh}^{(n)})a\| < \frac{1}{n}, \quad \forall g, h \in G, \quad \forall a \in F_n$$

$$\|r_g^{(n)} a - a r_g^{(n)}\| < \frac{1}{n}, \quad \forall g \in G, \quad \forall a \in F_n$$

$$\|(\sum_{g\in G} r_g^{(n)})a - a\| < \frac{1}{n}, \quad \forall a \in F_n.$$

For each $g \in G$ set $r_g := (r_g^{(n)})_n \in \ell_\infty(\mathbb{N}, A)$. Then by the inequalities above, for each $a \in \bigcup_{n\in\mathbb{N}} F_n$ we have $r_g r_h a = 0$ for all $g, h \in G$ with $g \neq h$, $(\alpha_g(r_h) - r_{gh})a = 0$ for all $g, h \in G$, $a r_g = r_g a$ for all $g \in G$, and $(\sum_{g\in G} r_g)a = a$. Since $\bigcup_{n\in\mathbb{N}} F_n$ is dense in the unit ball of B, the same equalities hold for every $a \in B$. Let $\pi\colon A^\infty \cap B' \to F(B, A)$ be quotient map. For each $g \in G$ put $p_g = \pi([r_g])$. Then $(p_g)_{g\in G}$ are mutually orthogonal projections that satisfy conditions (a) and (b) of (iv) of Proposition 1. Therefore, by the same proposition α has the Rokhlin property. □

The following is an example of a $\mathbb{Z}_2$-action with the Rokhlin property that does not have the multiplier Rokhlin property:

Example 1. Consider the space $X = \{\frac{1}{n} : n \in \mathbb{N}\}$ with the topology induced from $\mathbb{R}$. Let $\phi\colon X \to X$ be the map defined by $\phi(1/(2n-1)) = 1/(2n)$, and $\phi(1/(2n)) = 1/(2n-1)$, for all $n \in \mathbb{N}$. It is clear that ϕ is bijective, continuous, proper, its inverse is continuous, and that $\phi^2 = \mathrm{id}_X$. This implies that the map $\alpha\colon \mathrm{C}_0(X) \to \mathrm{C}_0(X)$ defined by $\alpha(f)(x) = f(\phi^{-1}(x))$ is a *-automorphism satisfying $\alpha^2 = \mathrm{id}_{\mathrm{C}_0(X)}$. Consider the action of $\mathbb{Z}_2$ on $\mathrm{C}_0(X)$ defined by this automorphism. Let us see that this action has the Rokhlin property. For each $n \in \mathbb{N}$, let $r_n^{(1)}, r_n^{(2)} \in \mathrm{C}_0(X)$ denote the

characteristic functions of the open subsets $\{1/(2k-1) : 1 \leq k \leq n\}$ and $\{1/(2k) : 1 \leq k \leq n\}$, respectively (note that these functions are continuous since the given sets are open). Then $r_n^{(1)}$ and $r_n^{(2)}$ are orthogonal, $\alpha(r_n^{(1)}) = r_n^{(2)}$, and $(r_n^{(1)} + r_n^{(2)})_{n\in\mathbb{N}}$ is an approximate unit for $\mathrm{C}_0(X)$. Since $\mathrm{C}_0(X)$ is commutative it follows that the given $\mathbb{Z}_2$-action has the Rokhlin property. It is clear that this action does not have the multiplier Rokhlin property since X can not be written as the union of two disjoint open sets X_1 and X_2 satisfying $\phi(X_1) = X_2$. (If $X = X_1 \cup X_2$ where X_1 and X_2 are disjoint and open, then either X_1 or X_2 contains a subset of X of the form $\{1/n : n > k\}$. In particular, it follows that $\phi(X_1) \neq X_2$.)

Let A be a C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A. Recall that a map $\omega\colon G \to \mathrm{U}(\mathrm{M}(A))$, where $\mathrm{U}(\mathrm{M}(A))$ denotes the unitary group of $\mathrm{M}(A)$, is said to be an *α-cocycle* if it satisfies $\omega_{gh} = \omega_g \alpha_g(\omega_h)$ for all $g, h \in G$. Let $\beta\colon G \to \mathrm{Aut}(A)$ be an action of G on A. The actions α and β are said to be *exterior equivalent* if there exists an α-cocycle ω such that $\beta_g(a) = \omega_g \alpha_g(a) \omega_g^*$ for all $g \in G$ and all $a \in A$.

Theorem 1. Let A and B be C*-algebras and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A. The following statements hold:

(i) If $\beta\colon G \to \mathrm{Aut}(B)$ is an action with the Rokhlin property, then the tensor action
$$\alpha \otimes_{\min} \beta\colon G \to \mathrm{Aut}(A \otimes_{\min} B)$$
has the Rokhlin property. Here $\otimes_{\min}$ denotes the minimal tensor product;

(ii) If α has the Rokhlin property and B is a G-invariant hereditary C*-subalgebra of A then the restriction of α to B has the Rokhlin property;

(iii) If α has the Rokhlin property and I is a G-invariant closed two-sided ideal of A then the action induced by α on A/I has the Rokhlin property;

(iv) If for any finite subset $F \subset A$ and any $\epsilon > 0$ there exist a C*-algebra B with an action $\beta\colon G \to \mathrm{Aut}(B)$ with the Rokhlin property, and an equivariant *-homomorphism $\phi\colon B \to A$ such that $\mathrm{dist}(a, \phi(B)) < \epsilon$ for all $a \in F$. Then α has the Rokhlin property;

(v) If (A, α) is a direct limit—with equivariant connecting *-homomorphisms—of a direct system $(A_\lambda, \alpha_\lambda)_{\lambda\in\Lambda}$, and every action $\alpha_\lambda\colon G \to \mathrm{Aut}(A_\lambda)$ has the Rokhlin property, then α has the Rokhlin property;

(vi) If α has the Rokhlin property and H is a subgroup of G then the restriction of α to H has the Rokhlin property.

(vii) If $\beta\colon G \to \mathrm{Aut}(A)$ is an action with the Rokhlin property and α and β are exterior equivalent then α has the Rokhlin property.

Proof. (i) Given that G is finite the tensor action $\alpha \otimes_{\min} \beta$ is well defined. Let us show that $\alpha \otimes \beta$ has the Rokhlin property. Since $A \otimes_{\min} B$ is the closure of the span of simple tensors, it is enough to show that for any $\epsilon > 0$ and any finite set $F \subset A \otimes_{\min} B$ consisting of simple tensors, there exist orthogonal positive contractions $(f_g)_{g\in G}$ in $A \otimes_{\min} B$ satisfying (i), (ii), and (iii) of Definition 2. Let $\epsilon > 0$ and
$$F = \{a_k \otimes b_k : a_k \in A,\, b_k \in B,\, 1 \leq k \leq n\}.$$
Choose an approximate unit $(u_i)_{i\in I}$ for A consisting of G invariant elements (this approximate unit exists by Lemma 1). Then there exists $i \in I$ such that
$$\|u_i a_k - a_k u_i\| < \frac{\epsilon}{2}, \qquad \|u_i a_k - a_k\| < \frac{\epsilon}{2}, \tag{3}$$
for $1 \leq k \leq n$. Choose Rokhlin elements $(r_g)_{g\in G} \subset B$ for β corresponding to the finite set $\{b_1, b_2, \cdots, b_n\}$ and to the number $\frac{\epsilon}{2}$. For each $g \in G$ put $f_g = u_i \otimes r_g$. Then the elements $(f_g)_{g\in G}$

are mutually orthogonal since the elements $(r_g)_{g\in G}$ are orthogonal. Now by the triangle inequality, the choice of $(r_g)_{g\in G}$, and the inequalities in (3) we get

$$\|(\alpha\otimes\beta)_g(f_h)-f_{gh}\|\leq\|u_i\otimes(\beta_g(r_h)-r_{gh})\|\leq\frac{\epsilon}{2}<\epsilon,$$

$$\|f_g(a_k\otimes b_k)-(a_k\otimes b_k)f_g\|\leq\|(u_ia_k-a_ku_i)\otimes r_gb_k\|+\|a_ku_i\otimes(r_gb_k-b_kr_g)\|<\frac{\epsilon}{2}+\frac{\epsilon}{2}=\epsilon,$$

$$\|(\sum_{g\in G}f_g)(a_k\otimes b_k)-(a_k\otimes b_k)\|\leq\|(u_ia_k-a_k)\otimes\sum_{g\in G}r_gb_k\|+\|a_k\otimes(\sum_{g\in G}r_gb_k-b_k)\|<\frac{\epsilon}{2}+\frac{\epsilon}{2}=\epsilon,$$

for $1\leq k\leq n$. We have shown that $(f_g)_{g\in G}$ are Rokhlin elements for β corresponding to the finite set F and the number ϵ. This implies that β has the Rokhlin property.

(ii) Let A, B, and α be as in the statement of (ii). Let F be a finite subset of B. Then by (ii) of Proposition 1 there exist orthogonal positive contractions $(r_g)_{g\in G}\subset A^\infty\cap F'$ satisfying $\alpha_g(r_h)=r_{gh}$ for all $g,h\in G$, and $(\sum_{g\in G}r_g)a=a$ for all $a\in F$. By Lemma 2 applied to the separable C*-subalgebra of B generated by the elements in F and to the finite sets $F_1=\{r_g:g\in G\}$, $F_2=\{0\}$, and $F_3=\{\sum_{g\in G}r_g\}$, there exists a positive contraction $d\in B^\infty\cap B'\subset A^\infty\cap B'$ such that $da=a$ for all $a\in F$, $dr_g=r_gd$ for all $g\in G$, and $(\sum_{g\in G}r_g)d=d$. For each $g\in G$ set $f_g=dr_gd$. Then $f_g\in B^\infty\cap B'$ since B is a hereditary C*-subalgebra of A. Also, it is straightforward to check that the elements $(f_g)_{g\in G}$ satisfy (a) and (b) of (ii) of Proposition 1. Therefore, by the equivalence of (i) and (ii) in the same proposition, the restriction of α to B has the Rokhlin property.

(iii) Let $\epsilon>0$ and let $F\subset A/I$ be a finite set. Let $\pi\colon A\to A/I$ denote the quotient map. Choose a finite set $\widetilde{F}\subset A$ such that $\pi(\widetilde{F})=F$. Choose Rokhlin elements $(r_g)_{g\in G}$ for $\widetilde{F}$ and $\epsilon>0$. Then it is straightforward that $(\phi(r_g))_{g\in G}$ are Rokhlin elements for F and ϵ.

(iv) Let A and α be as in the statement of the proposition. Let $F=\{a_1,a_2,\cdots a_n\}$ be a finite subset of A. Then by assumption there exist a C*-algebra B with an action $\beta\colon G\to\mathrm{Aut}(B)$ with the Rokhlin property, an equivariant *-homomorphism $\phi\colon B\to A$, and a finite subset $\widetilde{F}=\{b_1,b_2,\cdots b_n\}\subset B$ such that $\|a_i-\phi(b_i)\|<\frac{\epsilon}{3}$ for $1\leq i\leq n$. Since β has the Rokhlin property, there exist orthogonal positive contractions $(r_g)_{g\in G}\subset B$ such that

$$\|\alpha_g(r_h)-r_{gh}\|<\epsilon,\quad\|r_gb_i-b_ir_g\|<\frac{\epsilon}{3},\quad\|(\sum_{g\in G}r_g)b_i-b_i\|<\frac{\epsilon}{3},$$

for $g,h\in G$ and $1\leq i\leq n$. For each $g\in G$ set $f_g:=\phi(r_g)$. Then $(f_g)_{g\in G}$ are orthogonal positive contractions. Using that ϕ is equivariant we get

$$\|\beta_g(f_h)-f_{gh}\|=\|\phi(\alpha_g(r_h)-r_{gh})\|\leq\|\alpha_g(r_h)-r_{gh}\|<\epsilon.$$

Also, using the triangle inequality we get

$$\|f_ga_i-a_if_g\|\leq\|\phi(r_gb_i-b_ir_g)\|+\|f_g(a_i-\phi(b_i))\|+\|(a_i-\phi(b_i))f_g\|<\frac{\epsilon}{3}+\frac{\epsilon}{3}+\frac{\epsilon}{3}=\epsilon,$$

$$\|(\sum_{g\in G}f_g)a_i-a_i\|\leq\|\phi((\sum_{g\in G}r_g)b_i-b_i)\|+\|(\sum_{g\in G}f_g)(a_i-\phi(b_i))\|+\|a_i-\phi(b_i)\|<\frac{\epsilon}{3}+\frac{\epsilon}{3}+\frac{\epsilon}{3}=\epsilon,$$

for $g\in G$ and $1\leq i\leq n$. Hence, $(f_g)_{g\in G}$ are Rokhlin elements for α, F and ϵ. It follows that β has the Rokhlin property.

(v) This is a particular case of (iv).

(vi) Let H be a subgroup of G and let n denote the number of elements of G/H. Let $F\subset A$ and let $\epsilon>0$. Choose a family K of representatives of H in G and Rokhlin elements $(r_g)_{g\in G}$ for α, F, and ϵ/n. Then it is easy to check using the triangle inequality that $(f_h)_{h\in H}$, with $f_h=\sum_{g\in K}f_{hg}$ for all $h\in H$, are Rokhlin elements for the restriction of α to H, the finite set F, and the number $\epsilon>0$.

(vii) Let $\alpha, \beta\colon G \to \mathrm{Aut}(A)$ be actions of G on A that are exterior equivalent and suppose that α has the Rokhlin property. Let $\omega\colon G \to \mathrm{U}(\mathrm{M}(A))$ be the α-cocycle implementing the equivalence. Let F be a finite subset of A. Using Lemma 1 choose an approximate unit $(u_i)_{i\in I} \subset A$ for A that is quasicentral in $\mathrm{M}(A)$ and that satisfies $\alpha_g(u_i) = u_i$ for all $g \in G$ and $i \in I$. Then for each $n \in \mathbb{N}$ there exists $i_n \in I$ such that

$$\|u_{i_n} a - a\| < \frac{1}{n}, \quad \forall a \in F,$$
$$\|u_{i_n} w_g - w_g u_{i_n}\| < \frac{1}{n}, \quad \forall g \in G.$$

For each $n \in \mathbb{N}$ choose Rokhlin elements $(r_g^{(n)})_{g\in G} \subset A$ for the the finite set $F_n = F \cup \{u_{i_n} w_g : g \in G\}$ and the number $\frac{1}{n}$. Consider the elements $u = [(u_n)_n], r_g = [(r_g^{(n)})] \in A^\infty$. Then it follows that

$$ua = au = a, \qquad uw_g = w_g u, \tag{4}$$
$$\alpha_g(r_h) = r_{gh}, \qquad r_g a = a r_g, \quad r_g u w_g = u w_g r_g, \tag{5}$$
$$(\textstyle\sum_{g\in G} r_g) a = a, \quad (\sum_{g\in G} r_g) u w_h = u w_h, \tag{6}$$

for all $a \in F$ and $g, h \in G$. Note that by the third equality in (5) we have

$$u^2 r_g = (uw_g)(uw_g)^* r_g = (uw_g) r_g (uw_g)^* = r_g (uw_g)(uw_g)^* = r_g u^2,$$

for all $g \in G$. This implies that $f(u^2) r_g = r_g f(u^2)$ for every $f \in \mathrm{C}_0((0, \|u\|], \mathbb{R}_+)$, and in particular for square root function. Therefore, $u r_g = r_g u$ for all $g \in G$. For each $g \in G$ set $f_g := u r_g u$. Then $f_g = u^2 r_g = r_g u^2$. In addition, by the equalities in (4), (5), and (6) we get

$$\beta_g(f_h) = w_g \alpha_g(u r_h u) w_g^* = w_g u r_{gh} u w_g^* = u w_g r_{gh} (uw_g)^* = r_{gh} (uw_g)(uw_g)^* = r_{gh} u^2 = f_{gh},$$

for all $g, h \in G$,

$$f_g f_h = (u r_g u)(u r_h u) = u^2 r_g r_h u^2 = 0,$$

for all $g, h \in G$ with $g \neq h$,

$$f_g a = u r_g u a = u r_g a u = a u r_g u = a f_g,$$

for all $g \in G$ and $a \in F$, and

$$(\sum_{g\in G} f_g) a = \sum_{g\in G} u r_g u a = u (\sum_{g\in G} r_g) a = ua = a,$$

for all $a \in F$. This shows that $(f_g)_{g\in G}$ are elements of $A^\infty \cap F'$ that satisfy conditions (a) and (b) of (ii) of Proposition 1 for the action β and the finite set F. Therefore, it follows by the same proposition that β has the Rokhlin property. □

Recall that an action $\alpha\colon G \to \mathrm{Aut}(A)$ is said to be *pointwise outer* if, for $g \in G \setminus \{e\}$, the automorphism α_g is outer, that is, not of the form $a \mapsto \mathrm{Ad}(u)(a) = uau^*$ for some unitary u in the multiplier algebra $\mathrm{M}(A)$ of A.

Proposition 2. Let A be C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A with the Rokhlin property. Then α is pointwise outer.

Proof. Suppose that there exist $g_0 \in G \setminus \{e\}$ and a unitary u in the multiplier algebra $\mathrm{M}(A)$ of A such that $\alpha_{g_0}(a) = uau^*$ for all $a \in A$. Let b be a nonzero element of the fixed-point algebra

A^α. By (ii) of Proposition 1 applied to the finite set $F = \{b, bu^*\}$, there exist orthogonal positive contractions $(r_g)_{g\in G} \subset A^\infty$ such that

$$\alpha_g(r_h) = r_{g,h}, \quad r_g(bu^*) = (bu^*)r_g, \quad r_g b = br_g \quad (\sum_{g\in G} r_g)b = b,$$

for all $g, h \in G$. Note that $r_e b \neq 0$ since $\sum_{g\in G} \alpha_g(r_e b) = (\sum_{g\in G} r_g)b = b \neq 0$. Also, we have $(r_{g_0}b)(br_e) = (br_e)(r_{g_0}b) = 0$ and

$$ur_e bu^* = u(bu^*)r_e = \alpha_{g_0}(b)r_e = br_e.$$

Hence,

$$0 = \|\alpha_{g_0}(r_e b) - ur_e bu^*\| = \|r_{g_0}b - br_e\| = \max(\|r_{g_0}b\|, \|br_e\|) \neq 0,$$

which is a contradiction. Therefore α_g is outer for all $g \in G \setminus \{e\}$. □

Let A be a C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A. Let us denote by $\mathrm{Ideal}(A)$ the set of closed two-sided ideals of A, by $\mathrm{Ideal}(A)^G \subset \mathrm{Ideal}(A)$ the set of G-invariant ideals, and by $\mathrm{Ideal}(a)$ the closed two-sided ideal of A generated by $a \in A$. The following result is essentially [38, Theorem 1.25].

Proposition 3. Let A be C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A with the Rokhlin property. Then the map

$$\begin{aligned} \mathrm{Ideal}(A \rtimes_\alpha G) &\longrightarrow \mathrm{Ideal}(A)^G, \\ I &\longmapsto I \cap A, \end{aligned}$$

is a bijection. In particular, every ideal of $J \subset A \rtimes_\alpha G$ has the form $I \rtimes_{\alpha_{(\cdot)|_I}} G$ for some G-invariant ideal I of A.

Proof. The map above is clearly onto since $(I \rtimes_\alpha G) \cap A = I$ for all $I \in \mathrm{Ideal}(A)^G$. Let us show that it is one-to-one. Let $(u_g)_{g\in G} \subset \mathrm{M}(A \rtimes_\alpha G)$ be the canonical unitaries implementing the action α. By [38, Theorem 1.10(ii)] it is enough to show that for $x \in A \rtimes_\alpha G$, with $x = \sum_{g\in G} a_g u_g$, one has $a_e \in \mathrm{Ideal}(x)$.

Let $x = \sum_{g\in G} a_g u_g$ be an element of $A \rtimes_\alpha G$, with $a_g \in A$ for all $g \in G$. Choose Rokhlin elements $(r_g^{(n)})_{g\in G}$, with $n \in \mathbb{N}$, associated to the finite set $F = \{a_g : g \in G\}$ and to the numbers $\frac{1}{n}$, with $n \in \mathbb{N}$. For each $g \in G$ set $r_g = (r_g^{(n)})_n$. Then $r_g \in A^\infty \cap F' \subset (A \rtimes_\alpha G)^\infty \cap F'$ for all $g \in G$, $\alpha_g(r_h) = r_{gh}$ for all $g, h \in G$, and $\sum_{g\in G} r_g a_e = a_e$. It follows that

$$\sum_{h\in G} r_h x r_h = \sum_{g\in G}\sum_{h\in G} r_h a_g u_g r_h = \sum_{g\in G}\sum_{h\in G} \delta_{h,gh} r_h^2 a_g u_g = \sum_{h\in G} r_h^2 a_e = (\sum_{h\in G} r_h)^2 a_e = a_e.$$

Therefore, $\sum_{h\in G} r_h^{(n)} x r_h^{(n)} \to a_e$ as $n \to \infty$. This implies that $a_e \in \mathrm{Ideal}(x)$. □

Corollary 4. Let A be a simple C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A with the Rokhlin property. Then $A \rtimes_\alpha G$ is simple.

Proposition 4. Let A be C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A with the Rokhlin property. Then the crossed product C*-algebra $A \rtimes_\alpha G$ and the fixed point algebra A^G are strongly Morita equivalent.

Proof. This is a consequence of the discussion after [14, Lemma 3.1] (see also [27, Theorem 5.10]). □

Corollary 5. Let A be C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A with the Rokhlin property. Then $\mathrm{Cu}(A^G)$ and $\mathrm{Cu}(A \rtimes_\alpha G)$ are naturally isomorphic.

Proof. By Proposition 4 the crossed product C*-algebra $A \rtimes_\alpha G$ and the fixed point algebra A^G are strongly Morita equivalent. Hence, there exists a canonical C*-algebra B containing copies of $A \rtimes_\alpha G$ and A^G as full hereditary C*-subalgebras ([30, page 288]). It follows that $\mathrm{Cu}(A \rtimes_\alpha G) \cong \mathrm{Cu}(B) \cong \mathrm{Cu}(A^G)$. □

4. Local approximation of $A \rtimes_\alpha G$

In this section we show that the crossed product C*-algebra by an action of a finite group on a C*-algebra with the Rokhlin property can be locally approximated by matrix algebras over hereditary C*-subalgebras of the given C*-algebra.

Let $(f_{i,j})_{i,j=1}^d$ denote the set of matrix units of $\mathrm{M}_d(\mathbb{C})$. The following result is well known. We include the proof for the convenience of the reader.

Lemma 7. Let A be a C*-algebra, let $\phi\colon \mathrm{C}_0(0,1] \otimes \mathrm{M}_d(\mathbb{C}) \to A$ be a *-homomorphism, and let $b = \phi(t \otimes 1_{\mathrm{M}_d})$ and $c = \phi(t \otimes f_{1,1})$. Then the map $\psi\colon \mathrm{M}_d(cAc) \to \overline{bAb}$ defined by

$$\psi(\sum_{i,j=1}^d (cx_{i,j}c) \otimes f_{i,j}) := \sum_{i,j=1}^d \phi(t \otimes f_{i,1}) x_{i,j} \phi(t \otimes f_{1,j}),$$

extends to a *-isomorphism $\widetilde{\psi}\colon \mathrm{M}_d(\overline{cAc}) \to \overline{bAb}$.

Proof. The map ψ is clearly additive. Also, using that $(f_{i,j})_{i,j=1}^d$ is a set of matrix units of $\mathrm{M}_d(\mathbb{C})$ it is easy to check that ψ is multiplicative and that it preserves the adjoint operation.

Now note that the matrix $A_n = \left(\delta_{i,1}\phi(t^{\frac{1}{n}} \otimes f_{i,j})\right)_{1 \le i,j \le d}$, where $\delta_{i,j}$ denotes Kronecker's delta function, satisfies

$$\|A_n\| = \|A_n^* A_n\|^{\frac{1}{2}} = \|\sum_{i=1}^d \phi(t^{\frac{2}{n}} \otimes f_{i,i})\|^{\frac{1}{2}} \le 1.$$

Hence, we have

$$\begin{aligned}
\|\psi(\sum_{i,j=1}^n (cx_{i,j}c) \otimes f_{i,j})\| &= \|\sum_{i,j=1}^n \phi(t \otimes f_{i,1}) x_{i,j} \phi(t \otimes f_{1,j})\| \\
&= \lim_{n\to\infty} \|\sum_{i,j=1}^n \phi(t^{\frac{1}{n}} \otimes f_{i,1}) c x_{i,j} c \phi(t^{\frac{1}{n}} \otimes f_{1,j})\| \\
&= \lim_{n\to\infty} \|A_n^* \left(cx_{i,j}c\right)_{i,j} A_n\| \\
&\le \|\left(cx_{i,j}c\right)_{i,j}\| \\
&= \|\sum_{i,j=1}^n (cx_{i,j}c) \otimes f_{i,j}\|,
\end{aligned}$$

and

$$\begin{aligned}\|\sum_{i,j=1}^{n}(cx_{i,j}c)\otimes f_{i,j}\| &= \|\left(cx_{i,j}c\right)_{i,j}\| \\ &= \lim_{n\to\infty}\|A_n^t\left(\phi(t\otimes f_{i,1})x_{i,j}\phi(t\otimes f_{1,j})\right)_{i,j}(A_n^t)^*\| \\ &= \lim_{n\to\infty}\|\left(\phi(t^{1+\frac{1}{n}}\otimes f_{1,1})x_{i,j}\phi(t^{1+\frac{1}{n}}\otimes f_{1,1})\right)_{i,j}\| \\ &= \lim_{n\to\infty}\|A_n(\sum_{i,j=1}^{d}\phi(t\otimes f_{i,1})x_{i,j}\phi(t\otimes f_{1,j}))A_n^*\| \\ &\leq \|\sum_{i,j=1}^{d}\phi(t\otimes f_{i,1})x_{i,j}\phi(t\otimes f_{1,j})\| \\ &= \|\psi(\sum_{i,j=1}^{n}(cx_{i,j}c)\otimes f_{i,j})\|,\end{aligned}$$

for all $x_{i,j}\in A$. Therefore, ψ is an isometry and in particular it is well defined. Now using that $\mathrm{M}_d(cAc)$ is dense in $\mathrm{M}_d(\overline{cAc})$ we can extend ψ by continuity to an isometric *-homomorphism $\widetilde{\psi}\colon \mathrm{M}_d(\overline{cAc})\to\overline{bAb}$. Note that

$$\widetilde{\psi}(c^3\otimes 1_{\mathrm{M}_d})=\psi(c^3\otimes 1_{\mathrm{M}_d})=\sum_{i=1}^{d}\phi(t^3\otimes f_{i,i})=b^3.$$

Using that $\widetilde{\psi}$ preserves functional calculus we get $\widetilde{\psi}(c)=b$. This implies that $\widetilde{\psi}$ is surjective. Therefore, it is a *-isomorphism. □

Lemma 8. Let A be a C*-algebra and let $0<\epsilon<1$. Let $a,b\in A_+$ and $x\in A$ be such that

$$\|a-x^*x\|<\epsilon,\quad \|b-xx^*\|<\epsilon.$$

Then there exists $y\in A$ such that

$$(a-9\epsilon^{\frac{1}{2}})_+=y^*y,\quad yy^*\leq(b-\epsilon)_+,\quad \|y-x\|<16\epsilon^{\frac{1}{4}}.$$

Proof. Let a, b, and x be as in the statement of the lemma. Then by [32, Proposition 1] there exists $z\in A$ such that

$$(b-\epsilon)_+=zz^*,\quad z^*z\leq x^*x,\quad \|z-x\|<4\epsilon^{\frac{1}{2}}.$$

(A straightforward computation shows that [32, Proposition 1] holds for $C=4$.) In particular, we have

$$\|a-z^*z\|\leq\|a-x^*x\|+\|(x-z)^*x\|+\|z^*(x-z)\|<\epsilon+4\epsilon^{\frac{1}{2}}+4\epsilon^{\frac{1}{2}}\leq 9\epsilon^{\frac{1}{2}}.$$

By applying again [32, Proposition 1], this times to the elements a and z, we get an element $y\in A$ satisfying

$$(a-9\epsilon^{\frac{1}{2}})_+=y^*y,\quad yy^*\leq zz^*=(b-\epsilon)_+,\quad \|y-z\|<12\epsilon^{\frac{1}{4}}.$$

A simple computation now shows that $\|y-x\|<16\epsilon^{\frac{1}{4}}$. □

In order to prove the main theorem of this section we need the following lifting result. This is a slight refinement of [21, Theorem 10.2.2] for quotients of the form $\ell_\infty(\mathbb{N},A)/\mathrm{c}_0(\mathbb{N},A)$.

Proposition 5. Let A be a C*-algebra and let $0 < \epsilon < 1$. Given a *-homomorphism $\phi\colon \mathrm{M}_d(\mathrm{C}_0(0,1]) \to A^\infty$ and orthogonal positive contractions $h_1, h_2, \cdots, h_d \in \ell_\infty(\mathbb{N}, A)$ such that $[h_i] = \phi(t \otimes f_{i,i})$ for all $i \in \mathbb{N}$, there exists a *-homomorphism $\psi\colon \mathrm{M}_d(\mathrm{C}_0(0,1]) \to \ell_\infty(\mathbb{N}, A)$ that lifts ϕ and that satisfies

$$\psi(t \otimes f_{1,1}) \in \mathrm{C}^*(h_1), \quad \psi(t \otimes f_{i,i}) \leq h_i.$$

Proof. Let $\phi\colon \mathrm{M}_d(\mathrm{C}_0(0,1]) \to A^\infty$ and $h_1 = (h_{1,n}), h_2 = (h_{2,n}), \cdots, h_d = (h_{d,n}) \in \ell_\infty(\mathbb{N}, A)$ be as in the statement of the proposition. For each $1 \leq i \leq d$ let $x_i = (x_{i,n}) \in \ell_\infty(\mathbb{N}, A)$ be a lifting of $\phi(t \otimes f_{i,1})$. Then

$$\|h_{1,n}^2 - x_{i,n}^* x_{i,n}\| \to 0, \quad \|h_{i,n}^2 - x_{i,n} x_{i,n}^*\| \to 0,$$

as $n \to \infty$. By passing to a subsequence and relabeling we may assume that

$$\|h_{1,n}^2 - x_{i,n}^* x_{i,n}\| < \frac{1}{n}, \quad \|h_{i,n}^2 - x_{i,n} x_{i,n}^*\| < \frac{1}{n},$$

for all $1 \leq i \leq d$ and $n \in \mathbb{N}$. Using these inequalities and Lemma 8 we can find elements $y_{i,n}$ such that

$$y_{i,n}^* y_{i,n} = (h_{1,n} - 9\epsilon^{-\frac{1}{2}})_+, \quad y_{i,n} y_{i,n}^* \leq h_{i,n}^2, \quad \|y_{i,n} - x_{i,n}\| < 16\epsilon^{\frac{1}{4}}.$$

For each i set $y_i = (y_{i,n})_n$. Then it follows that y_i is an element in $\ell_\infty(\mathbb{N}, A)$ and that $[y_i] = [x_i] = \phi(t \otimes f_{i,1})$. Also, using the orthogonality of the elements h_i we get that $y_{i,n} y_{j,n} = 0$ for all $i, j \geq 2$, and that $y_{i,n}^* y_{j,n} = 0$ for all $i, j \geq 2$ with $i \neq j$. It follows now by [21, Proposition 3.3.1] that there exists a *-homomorphism $\psi\colon \mathrm{M}_d(\mathrm{C}_0(0,1]) \to \ell_\infty(\mathbb{N}, A)$ satisfying $\psi(t \otimes f_{i,1}) = y_i$ for all i. Therefore, we have

$$\psi(t \otimes f_{1,1}) = (y_1^* y_1)^{\frac{1}{2}} = ((h_1^2 - 9\epsilon^{-\frac{1}{2}})_+)^{\frac{1}{2}} \in \mathrm{C}^*(h_1),$$
$$\psi(t \otimes f_{i,i}) \leq (y_i y_i^*)^{\frac{1}{2}} \leq (h_i^2)^{\frac{1}{2}} = h_i,$$

for all i. (In order to obtain the last inequality we are using [24, Proposition 1.3.8].) □

Theorem 2. Let A be a C*-algebra and let G be a finite group. Let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action with the Rokhlin property. Then for any finite subset $F \subseteq A \rtimes_\alpha G$ and any $\epsilon > 0$ there exist a positive element $a \in A$ and a *-homomorphism $\varphi\colon \mathrm{M}_{|G|}(\overline{aAa}) \to A \rtimes_\alpha G$ such that $\mathrm{dist}(x, \mathrm{Im}(\varphi)) < \epsilon$ for all $x \in F$.

Proof. Let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action with the Rokhlin property and let $(u_g)_{g \in G} \subset \mathrm{M}(A \rtimes_\alpha G)$ be the canonical unitaries satisfying $\alpha_g(a) = u_g a u_g^*$ for all $a \in A$ and $g \in G$. Let F be a finite subset of $A \rtimes_\alpha G$. Since $A \rtimes_\alpha G$ is generated by the elements of the set $\{au_g : a \in A, g \in G\}$, we may assume that $F = \{au_g : a \in F_1, g \in G\}$, where F_1 is a finite subset of A such that $\alpha_g(F_1) = F_1$ for all $g \in G$.

By (ii) of Proposition 1 there exist orthogonal positive contractions $(r_g)_{g \in G} \subset A^\infty \cap F_1'$ satisfying

$$\alpha_g(r_h) = r_{gh}, \quad (\sum_{g \in G} r_g) b = b \sum_{g \in G} r_g = b, \tag{7}$$

for $g, h \in G$ and $b \in F_1$. Consider A^∞ as a C*-subalgebra of $(A \rtimes_\alpha G)^\infty$ and set $x_g := u_g r_e$ for $g \in G$. Then $x_g \in (A \rtimes_\alpha G)^\infty$, $\|x_g\| \leq 1$, and the following orthogonality relations hold

$$x_g x_h = 0 \quad \forall g, h \neq e, \quad x_g^* x_h = \delta_{g,h} r_e^2 \quad \forall g, h \in G.$$

This implies by [21, Proposition 3.3.1] that there exists a *-homomorphism $\phi\colon \mathrm{M}_{|G|}(\mathrm{C}_0(0,1]) \to (A \rtimes_\alpha G)^\infty$ such that $\phi(t \otimes f_{g,e}) = x_g$. Let $s_g = (s_{g,n}) \in \ell_\infty(\mathbb{N}, A)$, with $g \in G$, be orthogonal positive contractions that lift $(r_g)_{g \in G}$ (they exist by [21, Lemma 10.1.12]). Then by Proposition 5 the map ϕ can be lifted to a *-homomorphism $\psi = (\psi_n)\colon \mathrm{M}_{|G|}(\mathrm{C}_0(0,1]) \to \ell_\infty(\mathbb{N}, A \rtimes_\alpha G)$. Moreover, ψ

can be taken such that $\psi(t\otimes f_{e,e})\in \mathrm{C}^*(s_e)\subseteq \ell_\infty(\mathbb{N},A)$. Note that $\overline{\psi_n(t\otimes f_{e,e})A\psi_n(t\otimes f_{e,e})}$ is a hereditary C*-subalgebra of A. Also, using the equations in (7) we get

$$\begin{aligned}\lim_{n\to\infty}\psi_n(t\otimes 1_{\mathrm{M}_{|G|}})au_g&=\lim_{n\to\infty}\sum_{h\in G}\psi_n(t\otimes f_{h,h})au_g=au_g,\\ \lim_{n\to\infty}au_g\psi_n(t\otimes 1_{\mathrm{M}_{|G|}})&=\lim_{n\to\infty}au_g\sum_{h\in G}\psi_n(t\otimes f_{h,h})=\lim_{n\to\infty}a(\sum_{h\in G}\psi_n(t\otimes f_{h,h}))u_g=au_g,\end{aligned}\tag{8}$$

for $au_g\in F$.

Let us now use that $(r_g)_{g\in G}$ are orthogonal elements of $A^\infty\cap F_1'$ satisfying $\alpha_g(r_h)=r_{gh}$ for all $g,h\in G$. For each $au_k\in F$ we have

$$\begin{aligned}&\phi(t\otimes f_{e,g})(au_k)\phi(t\otimes f_{h,e})=x_g^*(au_k)x_h=r_eu_g^*au_ku_hr_e=u_g\alpha_g(r_e)au_ku_hr_e=u_g^*r_gau_{kh}r_e\\&=u_g^*r_ga\alpha_{kh}(r_e)u_{kh}=u_g^*r_gar_{kh}u_{kh}=u_g^*ar_gr_{kh}u_{kh}=\delta_{g,kh}u_g^*ar_g^2u_g=\delta_{g,kh}u_g^*au_g\alpha_{g^{-1}}(r_g^2)\\&=\delta_{g,kh}\alpha_{g^{-1}}(a)r_e^2=\delta_{g,kh}r_e\alpha_{g^{-1}}(a)r_e.\end{aligned}$$

Hence, since ψ is a lift of ϕ we get

$$\lim_{n\to\infty}\|\psi_n(t\otimes f_{e,g})(au_k)\psi_n(t\otimes f_{h,e})-\delta_{g,kh}\psi_n(t\otimes f_{e,e})\alpha_{g^{-1}}(a)\psi_n(t\otimes f_{e,e})\|=0.\tag{9}$$

Set $c_{g,n}=\psi_n(t\otimes f_{e,g})$, $d_n=\psi_n(t\otimes 1_{\mathrm{M}_{|G|}})$, $c_g=(c_{g,n})_{n\in\mathbb{N}}$, and $d=(d_n)_{n\in\mathbb{N}}$. Then by Lemma 7 there exist *-isomorphisms $\rho_n\colon \mathrm{M}_{|G|}(\overline{c_{e,n}(A\rtimes_\alpha G)c_{e,n}})\to\overline{d_n(A\rtimes_\alpha G)d_n}$ satisfying

$$\rho_n((c_{e,n}xc_{e,n})\otimes f_{g,h}):=\psi_n(t\otimes f_{g,e})x\psi_n(t\otimes f_{e,h})=c_{g,n}^*xc_{h,n}.$$

Denote by $\widetilde{\rho}_n\colon \mathrm{M}_{|G|}(\overline{c_{e,n}Ac_{e,n}})\to\overline{d_n(A\rtimes_\alpha G)d_n}$ the restriction of ρ_n to the C*-subalgebra $\mathrm{M}_{|G|}(\overline{c_{e,n}Ac_{e,n}})$.

Let $\epsilon>0$. By equations (8) and (9) we can choose n large enough such that

$$\begin{aligned}\|d_n^2(au_k)d_n^2-au_k\|&<\frac{\epsilon}{|G|^2+1},\\ \|c_{g,n}(au_k)c_{h,n}^*-\delta_{g,kh}c_{e,n}\alpha_{g^{-1}}(a)c_{e,n})\|&<\frac{\epsilon}{|G|^2+1},\end{aligned}$$

for all $au_k\in F$ and $g,h\in G$. It follows that

$$\begin{aligned}\|\widetilde{\rho}_n(\sum_{g\in G}(c_{e,n}\alpha_{g^{-1}}(a)c_{e,n})\otimes f_{g,k^{-1}g})-au_k\|&=\|\sum_{g,h\in G}\delta_{g,kh}\rho_n((c_{e,n}\alpha_{g^{-1}}(a)c_{e,n})\otimes f_{g,h})-au_k\|\\&\leq\frac{|G|^2}{|G|^2+1}\epsilon+\|\sum_{g,h\in G}\rho_n((c_{g,n}au_kc_{h,n}^*)\otimes f_{g,h})-au_k\|\\&=\frac{|G|^2}{|G|^2+1}\epsilon+\|\sum_{g,h\in G}(c_{g,n}^*c_{g,n}au_kc_{h,n}^*c_{h,n})-au_k\|\\&=\frac{|G|^2}{|G|^2+1}\epsilon+\|d_n^2au_kd_n^2-au_k\|\\&<\epsilon.\end{aligned}$$

This shows that $\mathrm{dist}(x,\mathrm{Im}(\widetilde{\rho}))<\epsilon$ for all $x\in F$. □

5. Permanence properties

In this section we show that several classes of C*-algebras are closed under crossed products by finite group actions with the Rokhlin property.

Lemma 9. Let A and let B be C*-subalgebra of A. Let $0 < \epsilon < 1$. Suppose that $a \in A$ is a positive contraction and $b \in B$ is such that $\|b - a\| < \epsilon$. Then

$$\|b - (a^*a)^{\frac{1}{2}}\| < 2\epsilon.$$

Proof. By the triangle inequality we have

$$\|b^2 - a^*a\| \leq \|b(b-a))\| + \|(b-a)^*a\| \leq (1 + \|b\|)\epsilon < (2+\epsilon)\epsilon < 2\epsilon.$$

Hence, by [7, Lemma 1] we get

$$\|b - (a^*a)^{\frac{1}{2}}\| < \sqrt{2}\epsilon^{\frac{1}{2}} < 2\epsilon.$$

□

Definition 3. Let $\mathcal{C}$ be a class of C*-algebras. A C*-algebra A is called a *local $\mathcal{C}$-algebra* if for every finite subset $F \subset A$ and every $\epsilon > 0$ there exist a C*-algebra B in $\mathcal{C}$ and a *-homomorphism $\phi\colon B \to A$ such that $\operatorname{dist}(x, \phi(B)) < \epsilon$ for all $x \in A$. When A is a unital C*-algebra and the C*-algebras in $\mathcal{C}$ and the *-homomorphisms ϕ are unital the C*-algebra A is called *unital local $\mathcal{C}$-algebra*. We say that $\mathcal{C}$ is *closed under local approximation* if every local $\mathcal{C}$-algebra (separable if the C*-algebras in $\mathcal{C}$ are separable) belongs to $\mathcal{C}$.

Note that any C*-algebra that is a direct limit of C*-algebras in $\mathcal{C}$ is a local $\mathcal{C}$-algebra. Also, if $\widetilde{\mathcal{C}}$ denotes the class of C*-algebras consisting of the unitization of the C*-algebras in $\mathcal{C}$ and $\widetilde{A}$ denotes the unitization of A, then $\widetilde{A}$ is a unital local $\widetilde{\mathcal{C}}$-algebra whenever A is a local $\mathcal{C}$-algebra. Similarly, if $\mathcal{C}^s$ denotes the class of C*-algebras consisting of the stabilization of the C*-algebras in $\mathcal{C}$, then $A \otimes \mathcal{K}$ is a local $\mathcal{C}^s$-algebra.

Lemma 10. Let $\mathcal{C}$ be a class of C*-algebras such that if B is a C*-algebra in $\mathcal{C}$ and J is a closed two-sided ideal of B then J is in $\mathcal{C}$. Let A be a local $\mathcal{C}$-algebra and let I be a closed two-sided ideal of A. Then I is a local $\mathcal{C}$-algebra.

Proof. Let $\mathcal{C}$, A, and I be as in the statement of the lemma. Let $\epsilon > 0$ and let $F = \{a_1, a_2, \cdots, a_n\}$ be a finite subset of I. Since every element of I is a linear combination of four positive contractions we may assume that the elements of F are positive contractions. Using Lemma 9 and that A is a local $\mathcal{C}$-algebra we may choose a C*-algebra $B \in \mathcal{C}$, a *-homomorphism $\phi\colon B \to A$ and positive elements $b_1, b_2, \cdots, b_n$ such that

$$\|a_i - \phi(b_i)\| < \epsilon,$$

for all $1 \leq i \leq n$. By [19, Lemma 2.2] there exist elements $d_i \in A$, with $1 \leq i \leq n$, such that

$$\phi((b_i - \epsilon)_+) = d_i^* a_i d_i \in I,$$

for all $1 \leq i \leq n$. Let J denote the preimage by ϕ of I. Then J is a closed two-sided ideal of B and thus, it belongs to $\mathcal{C}$. Also, by the previous equalities we have that $(b_i - \epsilon)_+ \in J$. The lemma now follows using that

$$\|a_i - \phi((b_i-\epsilon)_+)\| \leq \|a_i - \phi(b_i)\| + \|\phi(b_i) - \phi((b_i-\epsilon)_+)\| < \epsilon + \epsilon = 2\epsilon,$$

for all $1 \leq i \leq n$. □

Recall that a 1-dimensional noncommutative CW-complex ([10, Definition 2.4.1]), or shortly a 1-dimensional NCCW-complex, is a C*-algebra A that can be written as a pullback diagram of the form

$$\begin{array}{ccc} A & \longrightarrow & E \\ \downarrow & & \downarrow{\scriptstyle \phi} \\ \mathrm{C}([0,1], F) & \xrightarrow{\rho} & F \oplus F \end{array}$$

where E, F are finite dimensional C*-algebras, ϕ is an arbitrary $*$-homomorphism, and ρ is given by evaluation at 0 and 1. In contrast to [10, Definition 2.4.1], the maps ϕ and ρ are not assumed to be unital. Special cases of 1-dimensional NCCW-complexes are interval algebras (i.e., algebras of the form $\mathrm{C}([0,1], E)$, where E is a finite dimensional C*-algebra), and circle algebras (i.e., algebras of the form $\mathrm{C}(\mathbb{T}, E)$, where E is a finite dimensional C*-algebra). Direct limits of sequences of interval algebras are called AI-algebras whereas direct limits of sequences of circles algebras are called AT-algebras.

Proposition 6. The following classes of C*-algebras are closed under local approximation:

(i) Purely infinite C*-algebras;
(ii) C*-algebras of stable rank one;
(iii) C*-algebras of real rank zero;
(iv) C*-algebras of nuclear dimension at most n, where $n \in \mathbb{Z}_+$;
(v) Separable $\mathcal{D}$-stable C*-algebras, where $\mathcal{D}$ is a strongly self-absorbing C*-algebra;
(vi) Simple C*-algebras;
(vii) Separable C*-algebras whose quotients are stably projectionless;
(viii) Simple stably projectionless C*-algebras;
(ix) C*-algebras that are stably isomorphic to direct limits of sequences of C*-algebras in a class $\mathcal{S}$, where $\mathcal{S}$ is a class of finitely generated semiprojective C*-algebras that is closed under taking tensor products by matrix algebras over $\mathbb{C}$.
(x) Separable AF-algebras;
(xi) C*-algebras that are stable isomorphic to AI-algebras;
(xii) C*-algebras that are stable isomorphic AT-algebras;
(xiii) C*-algebras that can be expressed as a sequential direct limit of 1-dimensional NCCW-complexes;
(xiv) Separable C*-algebras with almost unperforated Cuntz semigroup;
(xv) Simple C*-algebras with strict comparison of positive elements.
(xvi) Separable C*-algebras whose closed two-sided ideals are nuclear and satisfy the Universal Coefficient Theorem.

Proof. In [19, Proposition 4.18], [31, Theorem 5.1], [4, Proposition 3.1], and [41, Proposition 2.3] it is shown that the classes of C*-algebras described in first four parts of the proposition are closed under taking arbitrary direct limits of sequences. The proofs that these classes of C*-algebras are closed under local approximation are essentially the same; thus, we will omit them.

(v) Let $\mathcal{C}$ be the class of $\mathcal{D}$-stable C*-algebras and let A be a local $\mathcal{C}$-algebra. Let $F \subset A$ and $G \subset \mathcal{D}$ be finite subsets and let $\epsilon > 0$. By Lemma 6 it is enough to show that there exists a completely positive contractive map $\rho\colon \mathcal{D} \to A$ satisfying (i), (ii), and (iii) of Lemma 6. Without loss of generality we may assume that the elements of G are contractions.

Given that A is a local $\mathcal{C}$-algebra, there exist a $\mathcal{D}$-stable C*-algebra B and a *-homomorphism $\phi\colon B \to A$ such that $\mathrm{dist}(a, \phi(B)) < \frac{\epsilon}{3}$, for all $a \in F$. Now we can choose a finite subset $\widetilde{F} \subset B$ satisfying $\mathrm{dist}(a, \phi(\widetilde{F})) < \frac{\epsilon}{3}$ for all $a \in F$. By Lemma 6 applied to the finite subsets $\widetilde{F} \subset B$ and $G \subset \mathcal{D}$, and to $\frac{\epsilon}{3}$, there exists a completely positive contractive map $\psi\colon \mathcal{D} \to B$ such that

$$\|b\psi(1_{\mathcal{D}}) - b\| < \frac{\epsilon}{3}, \quad \|b\psi(d) - \psi(d)b\| < \frac{\epsilon}{3}, \quad \|b(\phi(dd') - \phi(d)\phi(d'))\| < \frac{\epsilon}{3}, \tag{10}$$

for all $b \in \widetilde{F}$ and $d, d' \in G$. Put $\rho = \phi \circ \psi$. Then ρ is a completely positive contraction. For each $a \in F$ choose $b \in \widetilde{F}$ such that $\|a - \phi(b)\| < \frac{\epsilon}{3}$. Then using the triangle inequality and the

inequalities in (10) we get

$$\|a\rho(1_{\mathcal{D}}) - a\| \leq \|(a - \phi(b))\rho(1_{\mathcal{D}})\| + \|\phi(b - \psi(1_{\mathcal{D}})b)\| + \|\phi(b) - a\| < \epsilon,$$
$$\|a\rho - \rho(d)a\| \leq \|(a - \phi(b))\rho(d)\| + \|\phi(b\psi(d) - \psi(d)b)\| + \|\rho(d)(a - \phi(b))\| < \epsilon,$$
$$\|a(\rho(dd') - \rho(d)\rho(d'))\| \leq \|(a - \phi(b))(\rho(dd') - \rho(d)\rho(d'))\| + \|\phi(b(\psi(dd') - \psi(d)\psi(d')))\| < \epsilon,$$

for all $d, d' \in G$. This implies that ρ satisfies (i), (ii), and (iii) of Lemma 6. Therefore, A is $\mathcal{D}$-stable by Lemma 6.

(vi) Let $\mathcal{C}$ be the class of simple C*-algebras and let A be a local $\mathcal{C}$-algebra. Let $a_1, a_2 \in A$ be positive elements of norm one and let $0 < \epsilon < \frac{1}{2}$. Then there exist a simple C*-algebra B and a *-homomorphism $\phi\colon B \to A$ such that $\operatorname{dist}(a_i, \phi(B)) < \epsilon$ for $i = 1, 2$. Choose elements $b_1, b_2 \in B$ such that $\|a_i - \phi(b_i)\| < \epsilon$ for $i = 1, 2$. Note that by Lemma 9 we may assume that b_1 and b_2 are positive. Also, $(\phi(b_2) - \epsilon)_+$ is nonzero as $\epsilon < \frac{1}{2}$ and $\|a_2\| = 1$. By [19, Lemma 2.2] there exist $d_i \in A$, with $i = 1, 2$, such that

$$(a_1 - \epsilon)_+ = d_1^*\phi(b_1)d_1, \quad (\phi(b_2) - \epsilon)_+ = d_1^*a_2d_1.$$

This implies that $[(a_1 - \epsilon)_+] \leq [\phi(b_1)]$ and $[(\phi(b_2) - \epsilon)_+] \leq [a_2]$ in $\operatorname{Cu}(A)$. Since B is simple, $[b_1] \leq \sup_n n[(b_2 - \epsilon)_+]$ in $\operatorname{Cu}(B)$. Hence,

$$[(a_1 - \epsilon)_+] \leq [\phi(b_1)] = \operatorname{Cu}(\phi)([b_1]) \leq \sup_n n\operatorname{Cu}(\phi)([(b_2 - \epsilon)_+]) = \sup_n n[(\phi(b_2) - \epsilon)_+] \leq \sup_n n[a_2],$$

in $\operatorname{Cu}(A)$. This implies that $(a_1 - \epsilon)_+$ is in the closed two-sided ideal generated by a_2. Since this holds for every $\epsilon > 0$ and $a = \lim_{\epsilon \to 0}(a_1 - \epsilon)_+$, a_1 is also an element of this ideal. Using now that a_1 and a_2 are arbitrary we get that A is simple.

(vii) Let $\mathcal{C}$ be the class of separable C*-algebras B such that for every closed two-sided ideal I of B, $B/I \otimes \mathcal{K}$ is projectioneless. Let A be a local $\mathcal{C}$-algebra. By the remarks after Definition 3, and [36, Corollary 2.3 part (i)] we may assume that A and all the algebras in $\mathcal{C}$ are stable. Let I be an ideal of A and let $\pi\colon A \to A/I$ denote the quotient map. Suppose that there exists a projection $p \in A/I$ with $p \neq 0$. Choose a positive element $a \in A$ such that $\pi(a) = p$. Since A is a local $\mathcal{C}$-algebra, there exist a C*-algebra $B \in \mathcal{C}$, a *-homomorphism $\phi\colon B \to A$, a positive element $a' \in A$ (here we are also using Lemma 9) such that

$$\|\phi(b) - a\| < \frac{1}{4}.$$

By passing to the quotient of B by the kernel of ϕ and using that this quotient is projectionless we may assume that ϕ is injective. Put $J := \phi^{-1}(I)$. Then J is a closed two-sided ideal of B and ϕ induces an injective *-homomorphism $\overline{\phi}\colon B/J \to A/I$. Let b' be the image of b in B/J. Then using that $\|\phi(b) - a\| < \frac{1}{4}$ we get

$$\|\overline{\phi}(b') - p\| < \frac{1}{4}.$$

This implies that $\overline{\phi}(b')$ has gap in its spectrum by [34, Lemma 2.2.3]. Hence, by functional calculus $\overline{\phi}(B/J)$ contains a nonzero projection. Using that $\overline{\phi}$ is injective we get that B/J contains a nonzero projection, which contradicts the fact that B/J is projectionless. Therefore, A belongs to $\mathcal{C}$.

Note that the proof above (with some minor modifications) also works for the class of (not necessarily separable) simple stably projectionless C*-algebras. Hence, the class in (viii) is closed under local approximation.

(ix) Let $\mathcal{C}$ and $\mathcal{S}$ be the classes of C*-algebras described in (ix). Let A be a separable local $\mathcal{C}$-algebra. It suffices to show that $A \otimes \mathcal{K}$ can be expressed as a direct limit of a sequence of C*-algebras in $\mathcal{S}$. Since A is a local $\mathcal{C}$-algebra, $A \otimes \mathcal{K}$ is a local $\mathcal{S}^s$-algebra, where $\mathcal{S}^s = \{B \otimes \mathcal{K} : B \in \mathcal{S}\}$. It follows that $A \otimes \mathcal{K}$ can be locally approximated by C*-algebras of the form $\mathrm{M}_n(B)$, where $B \in \mathcal{S}$ and $n \in \mathbb{N}$ (note that by [21, Theorem 14.2.2] each of the algebras $\mathrm{M}_n(B)$ is semiprojective). This

implies that $A \otimes \mathcal{K}$ is a $\mathcal{S}$-algebra, given that $\mathcal{S}$ is closed by taking tensor products with matrix algebras over $\mathbb{C}$. It follows now by [10, Lemma 2.2.5] and [21, Lemma 15.2.2] that $A \otimes \mathcal{K}$ is a direct limit of a sequence of C*-algebras in $\mathcal{S}$.

Parts (x), (xi), (xii), and (xiii) are a consequence of (ix), by taking $\mathcal{S}$ to be the class of finite dimensional C*-algebras, the class of interval algebras, the class of circle algebras, and the class of 1-dimensional NCCW-complexes, respectively. Each of these algebras are semiprojective and finitely generated by [10, Lemma 2.4.3 and Theorem 6.2.2]. The fact that separable AF-algebras are closed under local approximation was first proved in [2, Theorem 2.2] using a different method.

(xiv) Let $\mathcal{C}$ be the class of separable C*-algebras with almost unperforated Cuntz semigroup. Let A be a local $\mathcal{C}$-algebra. Note that if $\mathcal{C}^s$ denotes the class of C*-algebras consisting of the stabilization of the algebras in $\mathcal{C}$, then $A \otimes \mathcal{K}$ is a local $\mathcal{C}^s$-algebra. Therefore, using that the Cuntz semigroup is a stable functor, we may assume that A and all the algebras in $\mathcal{C}$ are stable. Also, using that $\mathcal{C}$ is closed under taking quotients by Lemma 5, we may assume that the *-homomorphisms used in the approximation of A by algebras in $\mathcal{C}$ are injective (here we are using that all the algebras in $\mathcal{C}$ are separable; thus, all their closed two-sided ideal are σ-unital).

Since A is stable, it is enough to show that, for $a, b \in A$, if $(n+1)[a] \leq n[b]$ then $[a] \leq [b]$. Let $a, b \in A_+$ be such elements. Then it follows that

$$a \otimes 1_{\mathrm{M}_{n+1}} \precsim b \otimes 1_{\mathrm{M}_n}.$$

Let $\epsilon > 0$. Then by (ii) of Lemma 3 there exists $\delta > 0$ such that

$$(a-\epsilon)_+ \otimes 1_{\mathrm{M}_{n+1}} \precsim (b-\delta)_+ \otimes 1_{\mathrm{M}_n}.$$

By the definition of Cuntz subequivalence, there exists $d \in \mathrm{M}_{n+1}(A)$ such that

$$\|(a-\epsilon)_+ \otimes 1_{\mathrm{M}_{n+1}} - d^*((b-\delta)_+ \otimes 1_{\mathrm{M}_n})d\| < \epsilon.$$

Using now that A is a local $\mathcal{C}$-algebra, there exist a C*-algebra $B \in \mathcal{C}$, an injective *-homomorphism $\phi\colon B \to A$, elements $a', b' \in B_+$ (here we are also using Lemma 9) and an element $d' \in \mathrm{M}_{n+1}(B)$ such that

$$\begin{aligned} &\|a - \phi(a')\| < \epsilon, \quad \|b - \phi(b')\| < \delta, \\ &\|(\phi \otimes 1_{\mathrm{M}_{n+1}})\left((a'-\epsilon)_+ \otimes 1_{\mathrm{M}_{n+1}} - (d')^*((b'-\delta)_+ \otimes 1_{\mathrm{M}_n})d'\right)\| < \epsilon. \end{aligned} \tag{11}$$

Since ϕ is injective, it is isometric. Hence, by the last inequality we have

$$\|(a'-\epsilon)_+ \otimes 1_{\mathrm{M}_{n+1}} - (d')^*((b'-\delta)_+ \otimes 1_{\mathrm{M}_n})d'\| < \epsilon,$$

in $\mathrm{M}_{n+1}(B)$. It follows that

$$(a'-2\epsilon)_+ \otimes 1_{\mathrm{M}_{n+1}} \precsim (b'-\delta)_+ \otimes 1_{\mathrm{M}_n}$$

by (i) of Lemma 3. Hence, $(n+1)[(a'-2\epsilon)_+] \leq n[(b'-\delta)_+]$ in $\mathrm{Cu}(B)$. Using now that $\mathrm{Cu}(B)$ is almost unperforated we get $[(a'-2\epsilon)_+] \leq [(b'-\delta)_+]$. By the first two inequalities in (11) and (i) of Lemma 3 we get

$$(a-3\epsilon)_+ \precsim (\phi(a')-2\epsilon)_+, \quad (\phi(b')-\delta)_+ \precsim b.$$

Therefore,

$$[(a-3\epsilon)_+] \leq [(\phi(a')-2\epsilon)_+] = [\phi((a'-2\epsilon)_+)] \leq [\phi((b'-\delta)_+)] = [(\phi(b')-\delta)_+)] \leq [b].$$

Using now that $[a] = \sup_{\epsilon>0}[(a-3\epsilon)_+]$ we get $[a] \leq [b]$, as desired.

(xv) Let $\mathcal{C}$ be the class of simple C*-algebras with strict comparison of positive elements and let A be a local $\mathcal{C}$-algebra. Then by Lemma 4 every C*-algebra in $\mathcal{C}$ has almost unperforated Cuntz semigroup. Note now that the proof of (xii) also applies for the class $\mathcal{C}$ and the algebra A. The separability condition was only used to assure that the *-homomorphisms used in the local approximation of A are injective, and this is obviously true for simple C*-algebras. Hence, it follows

that A has almost unperforated Cuntz semigroup. Therefore, A has strict comparison of positive elements by Lemma 4.

(xvi) Let $\mathcal{C}$ be the class of separable C*-algebras whose closed two-sided ideals are nuclear and satisfy the Universal Coefficient Therem (shortly, the UCT). Let A be a local $\mathcal{C}$-algebra and let I be a closed two-sided ideal of A. Then I is a local $\mathcal{C}$-algebra by Lemma 10. It follows now by [9, Theorem 1.1] that I satisfies the UCT. That I is nuclear follows using Arveson's extension theorem and that I can be locally approximated by nuclear C*-algebras. □

Theorem 3. The following classes of C*-algebras are preserved under taking crossed products with actions of finite groups with the Rokhlin property:

(i) Purely infinite C*-algebras;
(ii) C*-algebras of stable rank one;
(iii) C*-algebras of real rank zero;
(iv) C*-algebras of nuclear dimension at most n, where $n \in \mathbb{Z}_+$;
(v) Separable $\mathcal{D}$-stable C*-algebras, where $\mathcal{D}$ is a strongly self-absorbing C*-algebra;
(vi) Simple C*-algebras;
(vii) Simple C*-algebras that are stably isomorphic to direct limits of sequences of C*-algebras in a class $\mathcal{S}$, where $\mathcal{S}$ is a class of finitely generated semiprojective C*-algebras that is closed under taking tensor products by matrix algebras over $\mathbb{C}$;
(viii) Separable AF-algebras;
(ix) Separable simple C*-algebras that are stably isomorphic to AI-algebras;
(x) Separable simple C*-algebras that are stably isomorphic to AT-algebras;
(xi) C*-algebras that are stably isomorphic to sequential direct limits of one-dimensional NCCW-complexes;
(xii) Separable C*-algebras whose quotients are stably projectionless;
(xiii) Simple stably projectionless C*-algebras;
(xiv) Separable C*-algebras with almost unperforated Cuntz semigroup;
(xv) Simple C*-algebras with strict comparison of positive elements.
(xvi) Separable C*-algebras whose closed two-sided ideals are nuclear and satisfy the Universal Coefficient Theorem.

Proof. Let A be a C*-algebra and let $\alpha\colon G \to \mathrm{Aut}(A)$ be an action of a finite group G on A with the Rokhlin property. Let $\mathcal{S}$ be a class of C*-algebras such that for $B \in \mathcal{S}$, $n \in \mathbb{N}$, and $b \in B_+$ one has $\mathrm{M}_n(A) \in \mathcal{S}$ and $\overline{bBb} \in \mathcal{S}$. Then by Theorem 2, if A is a local $\mathcal{S}$-algebra then so is the crossed product C*-algebra $A \rtimes_\alpha G$. Given that all the classes of C*-algebras described in the theorem are closed under local approximation by Proposition 6 and under tensor product by matrix algebras, it is sufficient to show that they are closed under taking σ-unital hereditary subalgebras. This clearly holds for the classes of C*-algebras in (vi), (vii), (ix), (x), (xii), and (xiii) (here we are using [3, Theorem 2.8]). For the classes in (i), (iii), (iv), (v), (viii), and (xvi) this follows by [19, Proposition 3.15(iii)], [4, Corollary 2.8], [41, Propostion 2.5], [40, Corollary 3.1], [11, Theorem 3.1], and [34, Propositions 2.1.2(ii) and 2.4.7(iii)], respectively. For the class of C*-algebras in (xiv) this follows using [3, Theorem 2.8] and [16, Proposition 2.3]. For the class of C*-algebras in (xv) this holds by (xiv) and [39, Lemma 6.1]. Finally, for the class of algebras in (xi) this follows by the following argument. Let B be a hereditary C*-subalgebra of a C*-algebra A that is stably isomorphic to a direct limit of a sequence of 1-dimensional NCCW-complexes. Then $B \otimes \mathcal{K}$ is isomorphic to an ideal of $A \otimes \mathcal{K}$ by [3, Theorem 2.8]. It follows that $B \otimes \mathcal{K}$ is the direct limit of a sequence of ideals of 1-dimensional NCCW-complexes. Each of these ideals is a direct limit of a sequence of 1-dimensional NCCW-complexes by [37, Lemma 3.11]. Hence, $B \otimes \mathcal{K}$ can be locally approximated by 1-dimensional NCCW-complexes. This implies by Proposition 6 that $B \otimes \mathcal{K}$ is

a direct limit of a sequence of such algebras. Therefore, B belongs to the class of C*-algebras in (xi). □

References

[1] R. Antoine, F. Perera, and L. Santiago, *Pullbacks, $C(X)$-algebras, and their Cuntz semigroup*, J. Funct. Anal. **260** (2011), no. 10, 2844–2880.

[2] O. Bratteli, *Inductive limits of finite dimensional C^*-algebras*, Trans. Amer. Math. Soc. **171** (1972), 195–234.

[3] L. G. Brown, *Stable isomorphism of hereditary subalgebras of C^*-algebras*, Pacific J. Math. **71** (1977), no. 2, 335–348.

[4] L. G. Brown and G. K. Pedersen, *C^*-algebras of real rank zero*, J. Funct. Anal. **99** (1991), no. 1, 131–149.

[5] B. Blackadar, L. Robert, A. P. Tikuisis, A. S. Toms, and W. Winter, *An algebraic approach to the radius of comparison*, Trans. Amer. Math. Soc. **364** (2012), no. 7, 3657–3674.

[6] A. Ciuperca, L. Robert, and L. Santiago, *The Cuntz semigroup of ideals and quotients and a generalized Kasparov stabilization theorem*, J. Operator Theory **64** (2010), no. 1, 155–169.

[7] A. Ciuperca, G. A. Elliott, and L. Santiago, *On inductive limits of type-I C^*-algebras with one-dimensional spectrum*, Int. Math. Res. Not. IMRN **11** (2011), 2577–2615.

[8] K. T. Coward, G. A. Elliott, and C. Ivanescu, *The Cuntz semigroup as an invariant for C*-algebras*, J. Reine Angew. Math. **623** (2008), 161–193.

[9] M. Dadarlat, *Some remarks on the universal coefficient theorem in KK-theory*, Operator algebras and mathematical physics (Constanţa, 2001), Theta, Bucharest, 2003, pp. 65–74.

[10] S. Eilers, T. A. Loring, and G. K. Pedersen, *Stability of anticommutation relations: an application of noncommutative CW complexes*, J. Reine Angew. Math. **499** (1998), 101–143.

[11] G. A. Elliott, *Automorphisms determined by multipliers on ideals of a C^*-algebra*, J. Functional Analysis **23** (1976), no. 1, 1–10.

[12] Th. Fack and O. Maréchal, *Sur la classification des symétries des C^*-algebres UHF*, Canad. J. Math. **31** (1979), no. 3, 496–523 (French).

[13] E. Gardella and L. Santiago, *Classification of actions of finite groups on C*-algebras*, preprint (2014).

[14] E. C. Gootman, A. J. Lazar, and C. Peligrad, *Spectra for compact group actions*, J. Operator Theory **31** (1994), no. 2, 381–399.

[15] R. H. Herman and V. F. R. Jones, *Models of finite group actions*, Math. Scand. **52** (1983), no. 2, 312–320.

[16] I. Hirshberg, M. Rørdam, and W. Winter, *$\mathcal{C}_0(X)$-algebras, stability and strongly self-absorbing C^*-algebras*, Math. Ann. **339** (2007), no. 3, 695–732.

[17] M. Izumi, *Finite group actions on C*-algebras with the Rohlin property. I*, Duke Math. J. **122** (2004), no. 2, 233–280.

[18] ———, *Finite group actions on C*-algebras with the Rohlin property. II*, Adv. Math. **184** (2004), no. 1, 119–160.

[19] E. Kirchberg and M. Rørdam, *Infinite non-simple C*-algebras: absorbing the Cuntz algebras $\mathcal{O}_\infty$*, Adv. Math. **167** (2002), no. 2, 195–264.

[20] A. Kishimoto, *On the fixed point algebra of a UHF algebra under a periodic automorphism of product type*, Publ. Res. Inst. Math. Sci. **13** (1977/78), no. 3, 777–791.

[21] T. A. Loring, *Lifting solutions to perturbing problems in C^*-algebras*, Fields Institute Monographs, vol. 8, American Mathematical Society, Providence, RI, 1997.

[22] N. Nawata, *Finite group actions on certain stably projectionless C*-algebras with the Rohlin property*, Preprint, arXiv:1308.0429 (2013).

[23] H. Osaka and N. C. Phillips, *Crossed products by finite group actions with the Rokhlin property*, Math. Z. **270** (2012), no. 1-2, 19–42.

[24] G. K. Pedersen, *C^*-algebras and their automorphism groups*, London Mathematical Society Monographs, vol. 14, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], London, 1979.

[25] ———, *Unitary extensions and polar decompositions in a C^*-algebra*, J. Operator Theory **17** (1987), no. 2, 357–364.

[26] N. C. Phillips, *Equivariant K-theory and freeness of group actions on C^*-algebras*, Lecture Notes in Mathematics, vol. 1274, Springer-Verlag, Berlin, 1987.

[27] ———, *Freeness of actions of finite groups on C^*-algebras*, Operator structures and dynamical systems, Contemp. Math., vol. 503, Amer. Math. Soc., Providence, RI, 2009, pp. 217–257.

[28] ———, *The tracial Rokhlin property for actions of finite groups on C^*-algebras*, Amer. J. Math. **133** (2011), no. 3, 581–636.

[29] ———, *The tracial Rokhlin property is generic*, Preprint, arXiv:1209.3859 (2012).

[30] M. A. Rieffel, *Morita equivalence for operator algebras*, Operator algebras and applications, Part I (Kingston, Ont., 1980), Proc. Sympos. Pure Math., vol. 38, Amer. Math. Soc., Providence, R.I., 1982, pp. 285–298.
[31] ———, *Dimension and stable rank in the K-theory of C^*-algebras*, Proc. London Math. Soc. (3) **46** (1983), no. 2, 301–333.
[32] L. Robert and L. Santiago, *Classification of C^*-homomorphisms from $C_0(0,1]$ to a C^*-algebra*, J. Funct. Anal. **258** (2010), no. 3, 869–892.
[33] L. Robert, *Classification of inductive limits of 1-dimensional NCCW complexes*, Adv. Math. **231** (2012), no. 5, 2802–2836.
[34] M. Rørdam, F. Larsen, and N. Laustsen, *An introduction to K-theory for C^*-algebras*, London Mathematical Society Student Texts, vol. 49, Cambridge University Press, Cambridge, 2000.
[35] M. Rørdam, *Classification of nuclear, simple C^*-algebras*, Classification of nuclear C^*-algebras. Entropy in operator algebras, Encyclopaedia Math. Sci., vol. 126, Springer, Berlin, 2002, pp. 1–145.
[36] ———, *Stable C^*-algebras*, Operator algebras and applications, Adv. Stud. Pure Math., vol. 38, Math. Soc. Japan, Tokyo, 2004, pp. 177–199.
[37] L. Santiago, *Reduction of the dimension of nuclear C*-algebras*, arXiv:1211.7159 (2012).
[38] A. Sierakowski, *The ideal structure of reduced crossed products*, Münster J. Math. **3** (2010), 237–261.
[39] A. S. Toms and A. Tikuisis, *On the structure of the Cuntz semigroup in (possibly) nonunital C*-algebras*, arXiv:1210.2235 (2012).
[40] A. S. Toms and W. Winter, *Strongly self-absorbing C*-algebras*, Trans. Amer. Math. Soc. **359** (2007), no. 8, 3999–4029.
[41] W. Winter and J. Zacharias, *The nuclear dimension of C^*-algebras*, Adv. Math. **224** (2010), no. 2, 461–498.
[42] W. Winter, *Strongly self-absorbing C^*-algebras are $\mathcal{Z}$-stable*, J. Noncommut. Geom. **5** (2011), no. 2, 253–264.

E-mail address: lsantiag@fields.toronto.edu

Fields Institute for Research in Mathematical Sciences, 222 College St, Toronto, ON M5T 3J1, Canada